

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
31 October 2002 (31.10.2002)

PCT


(10) International Publication Number  
**WO 02/085942 A2**

- (51) International Patent Classification<sup>7</sup>: **C07K 14/705**
- (21) International Application Number: PCT/CA02/00560
- (22) International Filing Date: 24 April 2002 (24.04.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/285,713 24 April 2001 (24.04.2001) US  
60/356,163 14 February 2002 (14.02.2002) US
- (71) Applicant (for all designated States except US): **MCGILL UNIVERSITY** [CA/CA]; 845 Sherbrooke Street West, Montréal, Quebec H3A 2T5 (CA).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **PHILIP, Anie** [CA/CA]; 4100 Kensington, Montréal, Quebec H4B 2V8 (CA). **TAM, Betty** [CA/CA]; 1475 Everest, Ville St. Laurent, Quebec H4R 2R7 (CA).
- (74) Agents: **DUBUC, J. et al.**; Goudreau Gage Dubuc, Stock Exchange Tower, Suite 3400, 800 Place Victoria, P.O. Box 242, Montreal, Quebec H4Z 1E9 (CA).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— without international search report and to be republished upon receipt of that report

[Continued on next page]

(54) Title: A 150 KDA ACCESSORY RECEPTOR FOR TGF- $\beta$ .

Inhibition of TGF $\beta$  signaling by r150 : Expression of r150 markedly decreases  
TGF $\beta$ -induced Smad2 phosphorylation.



(57) Abstract: The present invention relates to a TGF- $\beta$ 1 binding protein called r150. This protein has a GPI-anchor contained in r150 itself and not on a tightly associated protein and that it binds TGF- $\beta$ 1 with an affinity comparable to those of the signaling receptors. Furthermore, the released (soluble) form of this protein binds TGF- $\beta$ 1 independent of the types I and II receptors. Also, the soluble form inhibits the binding of TGF- $\beta$  to its receptor. In addition, evidence that r150 is released from the cell surface by an endogenous phospholipase C is provided. Also, the creation of a mutant human keratinocyte cell line with a defect in GPI synthesis which displays reduced expression of r150 is described. Our results using these mutant keratinocytes suggest that the membrane anchored form of r150 is a negative modulator of TGF-beta responses. These findings, taken together with the observation that r150 forms a heteromeric complex with the signaling receptors, suggest that this accessory receptor in either its membrane anchored or soluble form may antagonize TGF- $\beta$  responses in human keratinocytes. Experiments with mutants confirmed that TGF $\beta$ 1 activity can be modulated when the expression of the accessory receptor r150 is silenced. The complete nucleic acid and deduced amino acid sequences are now provided. The r150 cloned nucleic acid was used to study overexpression of r150. When r150 gene is overexpressed, TGF $\beta$  responses are increased. r150 and its derivatives or precursors (fragments, variants and nucleic acids encoding the same) will find a broad clinical utility, knowing that TGF $\beta$ 1 is an important cytokine.

WO 02/085942 A2



*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

**TITLE OF THE INVENTION**

A 150 kDa accessory receptor for TGF- $\beta$ 1.

**BACKGROUND OF THE INVENTION**

Transforming growth factor- $\beta$  (TGF- $\beta$ ) is a 25 kDa multi  
5 functional growth factor which plays a central role in the wound healing process  
(Roberts and Sporn, 1990; O'Kane and Ferguson, 1997). It is an important regulator  
of the immune response (Letterio and Roberts, 1998), angiogenesis,  
reepithelialization (Roberts and Sporn, 1990), extracellular matrix protein synthesis  
and remodeling (Peltonen et al, 1991; Yamamoto et al, 1994). During wound healing,  
10 re-epithelialization initiates the repair process which is characterized by recruitment  
of epidermal stem cells, keratinocyte proliferation and the formation of an epithelial  
tongue of migrating keratinocytes at the wound edge (Clark, 1996). TGF- $\beta$  is  
chemotactic to keratinocytes and induces the expression of integrins on the migrating  
epithelium (Hebda, 1988; Zambruno et al, 1995). In spite of its promigratory eff ct  
15 on keratinocytes, TGF- $\beta$  is a potent inhibitor to epithelial cell proliferation *in vitro*  
(Pietenpol et al, 1990) and *in vivo* (Glick et al, 1993). Targeted deletion of the TGF-  
 $\beta$ 1 gene in keratinocytes causes rapid progression to squamous cell carcinoma  
(Glick et al, 1994). In addition, the epidermis of transgenic mice expressing a  
dominant negative TGF- $\beta$  receptor exhibits a hyperplastic and hyperkeratotic  
20 phenotype (Wang et al, 1997). These results support the importance of proper  
expression of TGF- $\beta$  and regulation of its function in epidermal development and  
maintenance of epidermal homeostasis .

TGF- $\beta$  is a member of the TGF- $\beta$  superfamily which also  
include activins, inhibins, bone morphogenic proteins, growth differentiation factor 1  
25 (GDF-1) and glial-derived neurotropic growth factor (GDNF) (Kingsley, 1994).

There are three widely distributed TGF- $\beta$  receptors, type I, type  
II and type III, all of which have been cloned (Roberts and Sporn, 1990; Massague,  
1998). The types I and II receptors are both transmembrane serine/threonine kinases  
that are essential for TGF- $\beta$  signal transduction. The type III receptor, also known as

- 2 -

betaglycan, is a high molecular weight proteoglycan that is not required for signaling, but is believed to play a role in presenting the ligand to the type II receptor (Lopez-Casillas et al, 1993). Endoglin, is another TGF- $\beta$  receptor predominantly expressed on endothelial cells (Gougos and Letarte, 1990). According to the present model of

5 TGF- $\beta$  signal transduction, binding of TGF- $\beta$  to the type II receptor which is a constitutively active kinase, leads to the recruitment and phosphorylation of the type I receptor (Wrana et al, 1994). The activated type I kinase phosphorylates the central intracellular mediators of TGF- $\beta$  signalling known as the Smad proteins (Heldin et al, 1997). Smad proteins can be classified into three groups: the pathway restricted

10 Smads include the Smad2 and Smad3 which are phosphorylated by the type I receptor of TGF- $\beta$  or activin, while the Smads 1, 5 and 8 are implicated in BMP signalling. The phosphorylation of the pathway restricted Smads permits their interaction with the common Smad or Smad4 and this heteromeric complex then translocates into the nucleus where it regulates expression of target genes. Finally,

15 there inhibitory Smads which include the Smad 7 and Smad 6 prevent the phosphorylation of the R-Smads by the type I kinase. (Heldin et al, 1997, Massague, 1998; Wrana and Attisano, 2000)

In blood circulation, TGF- $\beta$ 1 is found bound to the carrier  $\alpha_2$  macroglobulin ( $\alpha_2$ M; Webb et al. 1998).  $\alpha_2$ M binds many other cytokines and

20 therefore lacks selectivity for TGF- $\beta$ 1.  $\alpha_2$ M polymorphism has been associated with Alzheimer's disease, which polymorphism is observed as a deletion in "the bait region" overlapping with TGF- $\beta$ 1 binding domain (Gonias et al. 2000 and Blacker et al 1998).

Although the types I and II receptors are central to TGF- $\beta$  signaling, it is possible that accessory receptors interacting with the signaling

25 receptors modify TGF- $\beta$  responses. For example, both endoglin and type III receptor which form heteromeric complexes with the type II receptor have been reported to modulate TGF- $\beta$  function. When overexpressed in myoblasts, endoglin inhibited while type III receptor enhanced TGF- $\beta$  responses (Letamendia et al, 1998). In

30 addition, endoglin was shown to antagonize TGF- $\beta$  mediated growth inhibition of human vascular endothelial cells (Li et al, 2000). Similarly, the newly identified type

I-like receptor BAMBI which associates with TGF- $\beta$  family receptors can inhibit signaling (Onichtchouk et al, 1999).

There are also a number of molecules that can impact TGF- $\beta$  signal transduction by interacting with one or both of the TGF- $\beta$  signaling receptors.

5 However, the exact physiological significance of many of these interactions are not clearly defined (for review, Massague, 1998). Three of these interacting proteins: the type II TGF- $\beta$  receptor interacting protein (TRIP-1) (Chen et al, 1995), B $\alpha$  ( $\alpha$  subunit of protein phosphatase A) (Griswold-Prenner, 1998), and serine-threonine kinase receptor associated protein (STRAP) (Datta et al, 1998) all contain the highly  
10 conserved tryptophan-aspartic acid (WD) repeats. WD domains are important in protein-protein interactions and cellular functions such as cell cycle progression and transmembrane signaling (Neer et al, 1994). TRIP-1 is phosphorylated through its interaction with the type II receptor kinase and exerts an inhibitory effect on TGF- $\beta$  induced PAI-1 gene transcription, but has no effect on TGF- $\beta$  mediated growth  
15 inhibition (Choy and Derynck, 1998). On the other hand, B $\alpha$  associates with the type I receptor and positively modulates TGF- $\beta$  action. Finally, STRAP can interact with both the type I and II receptors and when overexpressed, it exerts an inhibitory effect on TGF- $\beta$  mediated transcriptional activation. In addition, STRAP can also interact with the inhibitory Smad7, but not Smad6. STRAP's interaction with Smad7 exerts  
20 a stabilizing effect on Smad7's association with the activated type I kinase receptor which prevents Smad2/3's association and subsequent phosphorylation (Datta and Moses, 2000).

The immunophilin, FKBP12, interacts with the TGF- $\beta$  type I receptor and acts as a negative modulator of TGF- $\beta$  function (Wang et al, 1996). It  
25 can interact with unactivated type I receptor and functions to stabilize the quiescent receptor state by protecting phosphorylation sites in the GS domain. Upon ligand stimulation, heteromerization and subsequent phosphorylation of the GS domain by the TGF- $\beta$  type II kinase results in the release of FKBP12 (Chen et al, 1997; Huse et al, 1999). In contrast, the TGF- $\beta$  type I receptor associated protein-1 (TRAP-1)  
30 interacts only with the activated type I receptor kinase (Charnig et al, 1998). TRAP-1 is not phosphorylated by the type I kinase and TRAP-1's interaction is reported to

- 4 -

have an inhibitory effect on TGF- $\beta$  signaling . However, a recent report describes a different function for TRAP-1 (Wurthner et al, 2001). In this study, TRAP-1 was found to associate with inactive TGF- $\beta$  and activin receptor complexes and upon ligand stimulation, TRAP-1 is released. The conformationally altered TRAP-1 is then  
5 believed to associate and subsequently chaperone Smad4 to the activated Smad2. The  $\alpha$  subunit of ras farnesyl protein transferase (FNTA) preferentially interacts with the activated type I receptor and is considered a substrate because it is phosphorylated by the type I kinase and released thereafter (Kawabata et al, 1995). However the functional significance of this phenomenon remains unexplained. The  
10 accessory receptors, endoglin and type III receptor which form heteromeric complexes with the type II receptor have also been reported to modulate TGF- $\beta$  function. When overexpressed in myoblasts, endoglin inhibited while type III receptor enhanced TGF- $\beta$  responses (Letamendia et al, 1998). Glycosylphosphatidylinositol (GPI)-anchored proteins which lack transmembrane and cytoplasmic domains have  
15 also been shown to bind TGF- $\beta$ . These proteins have been identified on certain cell lines (Cheifetz and Massague, 1991), but the identity of these GPI-anchored proteins and the role they may play in TGF- $\beta$  signaling remain unknown. Recently, the present inventors reported the presence of GPI-anchored TGF- $\beta$  binding proteins on  
20 early passage human endometrial stromal cells (Dumont et al, 1995), human skin fibroblasts (Tam and Philip, 1998) and keratinocytes (Tam et al, 1998). On human keratinocytes, they identified a 150 kDa GPI-anchored TGF- $\beta$ 1 binding protein designated as r150 that can form a heteromeric complex with the types I and II TGF- $\beta$  receptors (Tam et al, 1998). In addition, they demonstrated that upon hydrolysis  
25 from the cell surface by phosphatidylinositol phospholipase C (PIPLC), the soluble form of r150, retains its ability to bind TGF- $\beta$ 1 in the absence of the types I and II receptors. In addition, it was demonstrated that the GPI anchor is contained in a protein with a molecular weight of 150 kDa (Tam et al, 2001). This novel GPI-anchored TGF- $\beta$ 1 binding protein, r150, has the potential to antagonize or potentiate  
30 TGF- $\beta$  action in keratinocytes. In the absence of the cDNA encoding r150, one way to examine the effect of r150's loss in TGF- $\beta$  signaling is to enzymatically release the binding protein by PIPLC treatment prior to testing for alterations in TGF- $\beta$  induced responses. However, the efficacy of exogenously added PIPLC is subject to

variability, being affected by pH, temperature, and acylation of GPI-anchored proteins (Shukla, 1982; Chen et al, 1998), thus results obtained may be difficult to interpret. In addition, GPI-anchored proteins that are released may get re-synthesized and re-inserted in the plasma membrane soon after PIPLC hydrolysis. Hence, as an alternative, we have created and isolated a keratinocyte cell line that is mutated in GPI anchor biosynthesis. These cells display a significant loss of r150 from their cell surface, thus allowing a comparative examination of TGF- $\beta$  mediated cellular responses in the GPI anchor deficient cell line versus the parental HaCat cells under stable experimental conditions

GPI-anchored proteins lack transmembrane and cytoplasmic domains, and are attached to the cell membrane via a lipid anchor in which the protein is covalently linked to a glycosyl phosphatidylinositol moiety. GPI-anchored proteins have been reported to have roles in intracellular sorting (Rodríguez-Boulan and Powell, 1992), in transmembrane signaling (Brown, 1993) and to associate with cholesterol and glycosphingolipid-rich membrane microdomains (Brown and London, 1998; Hooper, 1999). Also, the GPI anchor enables a protein to be selectively released from the membrane by phospholipases (Metz et al, 1994; Movahedi and Hooper, 1997). r150 was characterized as GPI-anchored, based on its sensitivity to phosphatidylinositol phospholipase C (PIPLC). However, it is important to rule out other possibilities, namely, (i) r150 is not itself GPI-anchored, but is tightly associated with a protein that is GPI-anchored, and therefore is susceptible to release by PIPLC; (ii) r150 is a complex of two lower molecular weight proteins which became inadvertently cross-linked during the affinity labeling procedure.

It is now demonstrated that the GPI-anchor is contained in r150 itself and not on a tightly associated protein and that it binds TGF- $\beta$ 1 with an affinity comparable to those of the signaling receptors. Furthermore, the released (soluble) form of this protein binds TGF- $\beta$ 1 independent of the types I and II receptors. Also, the soluble form inhibits the binding of TGF- $\beta$  to its receptor. In addition, we provide evidence that r150 is released from the cell surface by an endogenous phospholipase C. Also, a mutant human keratinocyte cell line with a defect in GPI synthesis was created, which display reduced expression of r150. The results using

- 6 -

these mutant keratinocytes suggest that the membrane anchored form of r150 is a negative modulator of TGF-beta responses. These findings, taken together with the observation that r150 forms a heteromeric complex with the signaling receptors, suggest that this accessory receptor in either its membrane anchored or soluble form and its down- or up-regulation may potentiate or antagonize TGF- $\beta$  responses in human keratinocytes, respectively.

The complete amino acid of a molecule named CD109 was recently disclosed as well as the nucleic acids encoding same (Lin et al. 2002). Sequences comparisons with those of r150 suggest that CD109 is a r150 variant. No definite role has been assigned to CD109 by Lin et al.

## SUMMARY OF THE INVENTION

This invention provides a molecule that binds TGF- $\beta$ 1 with a high level of selectivity. This molecule named r150 can be retrieved in a membrane anchored form or as a released free soluble form. Variants and parts of r150 which have the property to bind TGF- $\beta$ 1 are grouped under the name r150-like proteins or peptides. They include those defined in SEQ ID Nos: 2, 4, 8, 10 and 12. Their corresponding coding nucleic acids respectively defined in SEQ ID NOs: 1, 3, 5, 7, 9 and 11.

This invention provides for the use of a protein comprising any one of the following sequences in the making of a medication for inhibiting TGF- $\beta$ 1 activity in a biological tissue SEQ ID Nos: 2, 4, 6, 8, 10 and 12.

Also provided is the use of an antagonist to a protein comprising any one of the following sequences in the making of a medication for increasing TGF- $\beta$ 1 activity in a biological tissue: SEQ ID Nos: 2, 4, 6, 8, 10 and 12.

Also provided is the use of a nucleic acid encoding a protein comprising any one of the following sequences in the making of medication for inhibiting TGF- $\beta$ 1 activity in a biological tissue: SEQ ID Nos: 1, 3, 5, 7, 9 and 11.

Also provided is the use of a molecule which silences the expression of a nucleic



encoding a protein comprising any one of the following sequences in the making of medication for increasing TGF- $\beta$ 1 activity in a biological tissue: SEQ ID Nos: 1, 3, 5, 7, 9 and 11. Particularly, the silencing molecule is an antisense nucleic acid.

5 The present inventors being the first to elucidate the complete nucleic acid sequence of r150 and of its deduced amino acid sequence, this invention provides an isolated nucleic acid encoding a protein comprising any one of the following sequences : SEQ ID Nos: 2, 4, 6, 8, 10 and 12.

In a specific embodiment, the nucleic acid comprises any one of the following nucleotide sequences: SEQ ID No: 1, 3, 5, 7, 9 and 11.

10 The nucleic acid is particularly one comprising the nucleotide sequence defined in SEQ ID No: 1.

The above nucleic acids may include crude nucleic acids or recombinant vectors; namely expression vectors capable of governing transcription and translation of the crude nucleic acids inserted downstream to a  
15 promotor, are preferred tools for producing r150-like proteins.

Recombinant host cells which comprise the nucleic acids or the recombinant vector are other tools. The vectors are normally selected to comprise sequences compatible with the host's machinery. Intervening sequences located 5' and 3' with regard to the crude nucleic acids are adapted or selected by the  
20 skilled artisan desirous to produce a particular type of host cells. The signal peptide may be charged also for another one more appropriate for a given cell type.

There host cells may be domesticated and used in a method of producing a r150-like protein. Such a method comprises the steps of:

- 25
- growing a recombinant host cell in a culture medium supporting cell growth and expression of said nucleic acid;
  - recovering the protein from the culture medium or from the cell.

The nucleic acids may be antisense nucleic acids. They may be inserted in a recombinant vector, namely an expression vector and recombinant host cells which comprises such antisense nucleic acids can also be made.

It is further an object of this invention to provide a TGF- $\beta$ 1 binding reagent, which comprises a protein comprising any one of the following sequences: SEQ ID Nos 2, 4, 6, 8, 10 and 12.

Compositions of matter which comprise these reagents and a carrier are other objects of this invention.

The carrier may be a pharmaceutical carrier. Otherwise, it may be a solid support to which r150 is bound to immobilize TGF- $\beta$ 1.

## DESCRIPTION OF THE INVENTION

r150 is a TGF- $\beta$ 1 binding molecule. Its complete amino acid sequence as well as the nucleic acid sequence encoding same appear to have been first elucidated by the present inventors. Another group (Lin et al. 2002), using a very different approach (affinity binding to monoclonal antibodies) has found a blood cell surface antigen, which they called CD109. Sequence comparisons show that CD109 (SEQ ID Nos 5 and 6) comprises a 17 amino acid insertion at position 1218-1234 (51 nts). This addition accounts for the difference in amino acids number (1445 for CD109 versus 1428 for r150). Besides that, substitutions of nucleotides are noted: r150 amino acid thr<sup>1224</sup> is changed for a methionine (CD109 amino acid 1241). CD109 shows polymorphism at residue 703 (Schuh et al. 2002). A tyrosine or a serine represent different alleles of CD109. Such polymorphism would presumably exist for r150. It is possible that CD109 or r150 are responsible for the building of an immune response since allo antibodies are retrieved upon administration of CD109 isoforms. It may therefore be implied that an isoform compatible with the recipient subject's tissue may have to be selected as an administrable r150 active ingredient.

r 150 binds or sequesters TGF- $\beta$ 1, in its membrane anchored form as well as in its free soluble form (SEQ ID Nos: 4 and 8). As a result, TGF-  $\beta$ 1

availability is reduced. The effects induced by TGF-  $\beta$ 1 are therefore negatively modulated (or inhibited). Such inhibition may be desirable in condition where overproduction of TGF-  $\beta$ 1 leads to pathological states (cancer is a specific example of such pathology). On the contrary, in other occasions, increasing TGF-  $\beta$ 1 activity may be sought. For example, TGF-  $\beta$ 1 encourages tissue or organ graft success. Therefore silencing r150 would have for effect to increase TGF-  $\beta$ 1 availability and increase graft success.

r150 further appears to be related to  $\alpha_2$  macroglobulin ( $\alpha_2$ M) and since the TGF-  $\beta$  binding domain has been determined for  $\alpha_2$ M by Webb et al. (1998), the corresponding domain in r150 is presumed to be located a region corresponding to  $\alpha_2$  macroglobulin amino acids 666-706. These corresponds to r150 amino acids 651-683 (SEQ ID No: 10). Therefore, the r150 peptide having the sequence defined in SEQ ID NO: 10 is also contemplated as a TGF- $\beta$ 1 binding peptide within the scope of the invention; the nucleic acid encoding this peptide is as well.

Webb et al. (2000) even found the minimal  $\alpha_2$ M TGF- $\beta$ 1 binding sequence which appears to be delineated by amino acid 717 and 733. The corresponding stretch in r150 is found between amino acid residues 694 and 712 (SEQ ID No.12).

Gomas et al. (2000) reported that  $\alpha_2$ M gene polymorphism has been associated with Alzheimer's disease, which polymorphism is observed as a deletion in "the bait region" overlapping with TGF- $\beta$ 1 binding domain. It is envisageable that r150 could be useful to sequester and neutralize TGF-  $\beta$ 1 especially in diseases wherein  $\alpha_2$ M is deficient. Any portion of r150 or variants thereof that is capable of binding TGF-  $\beta$ 1 activity is intended to be used in the making of a medication or a method or composition for inhibiting TGF- $\beta$ 1 activity. This includes proteins or peptides comprising sequences defined in SEQ ID NOs. 2, 4, 6, 8, 10 and 12. These r150-like proteins or peptides would include any molecule having at least 50% homology with the above sequences. On the opposite, any molecule having an antagonistic activity to the above r150-like proteins or peptides would find a use in

- 10 -

th making of a medication or a method or a composition for increasing TGF- $\beta$ 1 activity.

5 Nucleic acids encoding the above r150-like proteins or peptides represent an alternative to the direct administration of proteins or peptides. Antisense nucleics would on the opposite silence the expression of r150-like proteins or peptides. All these nucleic acids include recombinants vectors, namely expression vectors, which are available and well known to the skilled artisan.

10 A very large body of literature describes diseases or disease models involving up and down regulation of TGF- $\beta$ 1 activity. Nowadays, TGF- $\beta$ 1 binding proteins decorin and an anti-TGF antibody are currently under clinical trials. The present r150-like proteins or peptides could represent a valuable and advantageous alternative to these molecules, because of their selectivity for TGF- $\beta$ 1 isoform, combined to their hydrosolubility.

15 Here is a non-exhaustive list of disease models where alteration of TGF- $\beta$  action has been shown to be of therapeutic benefit:

#### Cancer progression:

Note: TGF- $\beta$  has biphasic effects during tumorigenesis, acting early as a tumor suppressor, but later stimulating cancer progression.

#### (i) Suppression of tumor progression by TGF- $\beta$

20 Akhurst R.J. and Derynck R. (2001). TGF- $\beta$  signaling in cancer a double-edged sword. *TRENDS in Cell Biology* 11: S44 – S51.

Welch, Dr. et al (1990). Transforming growth factor-  $\beta$  stimulates mammary adenocarcinoma cell invasion and metastatic potential. Proc. Natl. Acad. Sci. USA 87: 7678-7682

5 Markowitz, S. et al. (1995) Inactivation of the type II TGF-  $\beta$  receptor in colon cancer cells with microsatellite instability. Science 268,1336-1338.

Massagué, J. et al. (2000) TGF-  $\beta$  signaling in growth control, cancer, and heritable disorders. Cell 103, 295-309.

(ii) Stimulation of tumor progression by TGF-  $\beta$

10 Hojo, M. et al. (1999) Cyclosporine induces cancer progression by a cell-autonomous mechanism. Nature 397, 530-534.

Yin, J.J. et al. (1999) TGF-  $\beta$  signaling blockade inhibits PTHrP secretion by breast cancer cells and bone metastases development. J. Clin. Invest. 103, 197-206.

15 Exogenous TGF-  $\beta$ 1 promotes wound healing where as inhibiting TGF-  $\beta$ 1 activity or enhancing TGF-  $\beta$ 3 activity reduces scarring in animal models

Roberts, A.B., and Sporn, M.B. (1996). Transforming growth factor beta. In: The molecular and cellular biology of wound repair (second edition), Clark, R.A.F. (ed), Plenum Press, New York, p275-308.

20 Mustoe, T.A., Pierce, G.F., Thomason, A., Gramates, P., Sporn, M.B., and Deuel, T.F. (1987). Accelerated healing of incisional wounds in rats induced by transforming growth factor-  $\beta$ . Science 237: 1333-1336.

Quaglino, D., Nann y, L.B., Ditesheim, J.A., and Davidson, J.M. (1991). Transforming growth factor- $\beta$  stimulates wound healing and modulates extracellular matrix gene expression in pig skin: incisional wound model. *J. Invest. Dermatol.* 97: 34-42.

- 5 O'Kane, S., and Ferguson, M.W.J. (1997). Transforming growth factor- $\beta$ s and wound healing. *Int. J. Biochem. Cell. Biol.* 29:63-78.

Shah, M., Foreman, D.M., and Ferguson, M.W.J. (1995). Neutralization of TGF- $\beta$ 1 and TGF- $\beta$ 2 or exogenous addition of TGF- $\beta$ 3 to cutaneous rat wounds reduces scarring. *J. Cell Science* 108: 985-1002.

- 10 Choi, B-M., Kwak, H-J., Jun, C-D., Park, S-D., Kim, K-Y., Kim, H-R., and Chung, H-T. (1996). Control of scarring in adult wounds using antisense transforming growth factor- $\beta$ 1 oligodeoxynucleotides. *Immunol. Cell Biol.* 74:144-150.

- 15 Blocking TGF- $\beta$ 1 overproduction reduce tissue fibrosis (pulmonary fibrosis, liver cirrhosis, glomerulonephritis, scleroderma and atherosclerosis).

Border et al, (1990). Suppression of experimental glomerulonephritis by antiserum against transforming growth factor beta 1. *Nature* 346 (6282): 371-374

- 20 Isaka Y., Brees D.K., Ikegaya K., Kaneda Y., Imai E., Noble N.A., Border W.A. (1996). Gene therapy by skeletal muscle expression of decorin prevents fibrotic disease in rat kidney. *Nat. Med.* 2 : 418-423.

Isaka Y., Akagi Y., Ando Y., Tsujie M., Sudo T., Ohno N., Border W.A., Noble N.A., Kaneda Y., Hori M., and Imai E. (1999). Gene therapy by transforming growth factor-beta receptor-IgG Fc chimera suppressed extracellular

matrix accumulation in experimental glomerulonephritis [see comments]. *Kidney Int.* 55:465-475.

Khaliq, N. and Greenberg AH (1991). The role of TGF- $\beta$  in pulmonary fibrosis. *Ciba Found Symp.* 157: 194-207.

- 5 Yamamoto, T., Takagawa S., Katayama I., and Nishioka K. (1999). Anti-Sclerotic effect of transforming growth factor -beta antibody in a mouse model of bleomycin-induced scleroderma. *Clin Immunol*, 92(1):6-13.

Gressner, A.M., Weiskirchen, R., Breitkopf K., and Dooley S. (2002). Roles of TGF-beta in hepatic fibrosis. *Front Biosci* (7): d793-807.

- 10 Sheppard D. (2001). Integrin-mediated activation of transforming growth factor-beta(1) in pulmonary fibrosis. *Chest* 120(1 Suppl):49S-53S.

McCaffrey, T.A. (2000). TGF- $\beta$ s and TGF- $\beta$  receptors in atherosclerosis. *Cytokine. Growth Factor Rev.* 11: 103-114.

- 15 TGF-beta has tissue protective effects (against ischemia reperfusion injury) in the heart, brain and kidney.

Lefer AM., Ma X-L, Weyrich AS, Scalia R. (1993). Mechanism of the cardioprotective effect of TGF- $\beta$ 1 in feline myocardial ischemia and reperfusion. *Proc. Natl. Acad. Sci. USA* 90: 1018-1022.

- 20 Lefer AM, Tsao P, Aoki N, Palladino MA. (1990). Mediation of cardioprotection by transforming growth factor- $\beta$ . *Science* 249: 61-64.

McNeill H, Williams C, Guan J, Dragunow M, Lawlor P, Sirimann E, Nikolics K,

- 14 -

Gluckman P. (1994). Neuronal rescue with transforming growth factor-beta 1 after hypoxic-ischaemic brain injury. *Neuroreport* 5: 901-904.

Mehta JL, Yang BC, Strates BS, Mehta P. (1999). Role of TGF-beta1 in platelet-mediated cardioprotection during ischemia-reperfusion in isolated  
5 rat hearts. *Growth Factors* 16: 179-190.

Recombinant hosts comprising the above nucleic acids or recombinant expression vectors can be used as a biological machinery in the production of the r150-like proteins or peptides. The elucidation of the nucleic acid sequence of r150 therefore leads to a method of producing these proteins or  
10 peptides by recombinant technology.

A variety of TGF- $\beta$ 1 binding reagents and compositions may be derived from the present invention.

First, peptides such as those defined in SEQ ID. Nos: 10 and 12 may be used as such as a TGF- $\beta$ 1 binding reagent. Larger molecules like those  
15 defined in SEQ ID Nos 2, 4, 6 and 8 could be conjugated (through their anchoring region) to a carrier. The carrier may take the form, for example, of a solid or semi-solid medium (beads, chromatography columns, plates, etc.), to immobilize TGF- $\beta$ 1. Pharmaceutical compositions would take any suitable form, depending on the selected route of administration. A r150-like protein or peptide (SEQ ID Nos: 2, 4, 8,  
20 8, 10 and 12) would be formulated with a pharmaceutically acceptable carrier. Doses equivalent to those used by intravenous route for decorin and/or the TGF-antibody can be produced.

Objects, advantages and features of the present invention will become more apparent upon reading of the following non restrictive description of preferred embodiments thereof, given by way of example only with reference to the  
25 accompanying drawings.

## BRIEF DESCRIPTION OF THE FIGURES



**Figure 1:** Affinity cross-link labeling of human neonatal keratinocytes with  $^{126}\text{I}$ -TGF- $\beta$ 1. Confluent monolayers were affinity labeled with 100 pM  $^{126}\text{I}$ -TGF- $\beta$ 1 in the absence or presence of unlabeled TGF- $\beta$ 1, - $\beta$ 2, or - $\beta$ 3. Solubilized cell extracts were analyzed by SDS-PAGE under non reducing conditions and autoradiography (a). Competition curves for r150 and the types I and II TGF- $\beta$  receptors were derived by densitometric analysis of a typical autoradiogram. The data for each binding complex are expressed as a percent of the value in control wells incubated with  $^{125}\text{I}$ -TGF- $\beta$ 1 alone and are plotted against the concentration of unlabeled TGF- $\beta$ 1 (b), - $\beta$ 2 (c), or - $\beta$ 3 (d). The autoradiogram and competition curves are representative of three different experiments.

**Figure 2:** Temperature induced phase separation in Triton X-114 of TGF- $\beta$  binding proteins on human neonatal keratinocytes. Keratinocytes affinity labeled with  $^{125}\text{I}$ -TGF- $\beta$ 1 were lysed in 1% Triton X-114. The Triton X-114 soluble material was incubated at 30°C for 10 minutes followed by centrifugation at room temperature to induce phase separation of the detergent rich phase and the aqueous detergent poor phase. A: An aliquot (20%) from each phase was precipitated with ethanol/acetone, and analyzed by SDS-PAGE under reducing conditions.  $^{126}\text{I}$ -TGF- $\beta$ 1 labeled proteins in the aqueous (Aq) and the detergent rich (Det) phases, representative of three different experiments, are shown. B: The remaining 80% of the detergent phase was utilized to determine the effect of PIPLC treatment on the partitioning of r150 in Triton X-114. The detergent phase was incubated in the absence (-) or presence (+) of 0.6 U/ml of PIPLC followed by temperature induced phase separation and ethanol/acetone precipitation as above, to distinguish between the hydrophilic and amphipathic forms of the proteins. Analysis of the  $^{126}\text{I}$ -TGF- $\beta$ 1 labeled proteins in the aqueous phases of PIPLC treated (+) and untreated (-) samples, by SDS-PAGE under reducing conditions are shown. The results shown are representative of two different experiments.

**Figure 3:** (a) Affinity labeling of soluble r150 with  $^{125}\text{I}$ -TGF- $\beta$ 1. To verify that the soluble form of the r150 can bind to TGF- $\beta$ 1, human keratinocytes (HaCaT) were left untreated (-) or treated with 0.6 U/ml of PIPLC (+). The GPI-anchored proteins released into the supernatant were concentrated and an aliquot

- 16 -

was affinity cross-link labeled with 150 pM of  $^{125}\text{I}$ -TGF- $\beta$ 1 in the absence or presence of excess unlabeled TGF- $\beta$ 1 and subjected to SDS-PAGE under reducing conditions. The result shown is representative of four different experiments. (b) Inhibition of  $^{125}\text{I}$ -TGF- $\beta$ 1 binding to TGF $\beta$  receptors by soluble r150. Confluent monolayers of HaCaT cells grown in T-25cm<sup>2</sup> culture flasks were left untreated or treated with 0.6 U/ml of PIPLC for 60 minutes at 37°C. The supernatants were collected and concentrated by Centricon. MvLu1 cells were affinity labeled with 50 pM  $^{125}\text{I}$ -TGF- $\beta$ 1 in the absence or presence of indicated doses of supernatants and  $^{125}\text{I}$ -TGF- $\beta$ 1 specifically bound was plotted as a function of the amount of supernatant used. The arbitrary unit of "1" is equivalent to a dose of supernatant from a T-25cm<sup>2</sup> flask (approximately  $1 \times 10^6$  cells). (c) Effect of anti-TGF- $\beta$ 1 on the inhibition of TGF- $\beta$ 1 binding to TGF- $\beta$  receptors. Confluent monolayers of MvLu1 cells were affinity labeled with 50 pM  $^{125}\text{I}$ -TGF- $\beta$ 1 in the absence (C), or presence of PIPLC treated supernatant: (+PIPLC - S), or PIPLC-S pretreated with non-immune rabbit IgG (15 $\mu$ g/ml), or PIPLC-S pretreated with anti-TGF- $\beta$ 1 antibody (15 $\mu$ g/ml). To demonstrate that the anti-TGF- $\beta$ 1 antibody effectively neutralizes TGF- $\beta$ , experiments were also performed with 100 pM of TGF- $\beta$ 1 (+ $\beta$ 1),  $\beta$ 1 pretreated with non-immune rabbit IgG (15 $\mu$ g/ml), or  $\beta$ 1 pretreated with anti-TGF- $\beta$ 1 antibody (15 $\mu$ g/ml). The values shown in (b) and (c) are the mean (+/- S.D.) of at least three to five different experiments.

**Figure 4: Identification of a GPI-anchor in r150.** Human neonatal keratinocytes were harvested and treated with PIPLC (as described in Materials and Methods). The supernatant containing the GPI-anchored proteins were purified using a TGF- $\beta$ 1 affinity column (see Materials and Methods for details). After the addition of the sample to the column, 0.5 ml fractions were collected during washing and elution. (a) An aliquot from each fraction was affinity labeled with 150 pM of  $^{125}\text{I}$ -TGF- $\beta$ 1 in the absence or presence of excess unlabeled TGF- $\beta$ 1 and samples were analyzed by SDS-PAGE under reducing conditions. Only fraction 21 demonstrated an affinity labeled protein at 150 kDa while in adjacent fractions no 150 kDa band was detectable. Affinity labeling pattern obtained for fraction 21 and fraction 18 are shown. (b) Selected fractions were subjected to SDS-PAGE and transferred to a nitrocellulose membrane, and the samples were immunoblotted with an anti-CRD antibody (Oxford GlycoSystems). A 150 kDa protein was detected in

fraction 21 but not in adjacent fractions (fraction 18). Immunoblotting with the anti-CRD antibody was performed twice and the affinity cross-link labeling experiments of soluble r150 was done at least three times.

**Figure 5:** Immunoprecipitation of affinity labeled TGF- $\beta$  binding complexes on human neonatal keratinocytes with the anti-CRD antibodies. Keratinocytes not treated with PIPLC were affinity labeled with 100 pM  $^{125}$ I-TGF- $\beta$ 1 (a & b) or  $^{126}$ I-TGF- $\beta$ 2 (c) and were not immunoprecipitated (nip) or subjected to immunoprecipitation with an anti-CRD antibody against trypanosomal sVSG (Oxford GlycoSystems) (a & c), or with an anti-CRD antibody against porcine membrane dipeptidase (Broomfield and Hooper, 1993) (c). In the lane marked anti-CRD+peptide the immunoprecipitation was carried out using the anti-CRD antibody which was preincubated with PIPLC treated membrane dipeptidase. Immune complexes were subjected to SDS-PAGE under reducing conditions and analyzed by autoradiography. The results shown are representative of at least four to five experiments.

**Figure 6 A:** Expression of CD59 is decreased in keratinocytes mutated in GPI anchor biosynthesis. A representative histogram of the expression of CD59 in a GPI anchor deficient clone, GPI M (white columns) and parental HaCat cells (black columns) as assessed by flow cytometry using an anti-CD59-FITC labeled antibody. The immunofluorescence intensity of CD59 in the GPI anchor deficient cells was approximately 50% to that of the HaCat cells. The control performed in the absence of antibody is also included (dotted peak). Flow cytometry was performed at least three times.

**Figure 6B:** Cell surface expression of r150 is markedly decreased in GPI anchor mutated cells. Confluent monolayers of HaCat and GPI M cells were affinity labeled with 100 pM  $^{126}$ I-TGF- $\beta$ 1. Solubilized cell extracts were analyzed on SDS-PAGE under reducing conditions.

**Figure 7 A:** Doubling time curves of HaCat and GPI anchor mutated cells. HaCat and GPI M cells were seeded at  $8.0 \times 10^5$  cells in 60 mm dishes

in duplicate and the cell number for each was determined at 14, 24, 28, 32, 36, 48, 52 and 72 hours using a hemacytometer.

**Figures 7 B and C:** The cellular morphology of GPI anchor mutated cells is identical to that of HaCat cells. Microscopic representation (10x magnification) of HaCat (Fig. 7B) or GPI M cells (Fig. 7C) under normal culture conditions.

**Figures 8 A and B:** Enhanced TGF- $\beta$  stimulated transcriptional response in GPI anchor mutated cells. HaCat and GPI M cells were transiently transfected with 1  $\mu$ g of p3TP-Lux reporter gene construct and were left untreated or treated under serum free conditions with the indicated concentrations of TGF- $\beta$ 1 for 4hrs (Fig. 8A) or 16 hrs (Fig. 8 B). The luciferase activity was normalized to  $\beta$ -galactosidase activity expressed from a co-transfected CMV $\beta$ gal plasmid. The data is representative of at least three independent experiments. Error bars represent standard deviation.

**Figures 9 A and B:** GPI anchor mutated cells display enhanced responses at low doses of TGF- $\beta$ 1. HaCat and GPI M cells were left untreated or treated with the indicated doses of TGF- $\beta$ 1 (Fig. 9 A) and TGF- $\beta$ 2 (Fig. 9 B) under serum free conditions for 20 minutes. Immunoblotting was performed using a rabbit polyclonal antibody specific to the phosphorylated form of Smad2. Immunoblotting was repeated with an anti-Smad2 antibody that recognizes total Smad2 (unphosphorylated and phosphorylated forms) or an anti-STAT3 antibody to demonstrate equal protein loading. This data is representative of three different experiments.

**Figures 10 A, B and C:** Elevated Smad2 phosphorylation is sustained in GPI anchor mutated cells. HaCat and two GPI anchor deficient clones GPI M (Fig. 10 A) and GPI M1 (Fig. 10 B) or a keratinocyte clone that was not GPI anchor mutated, GPI NM (Fig. 10 C) were treated with 100 pM TGF- $\beta$ 1 for the indicated times. The control lane (-) received no TGF- $\beta$ 1 treatment. Immunoblotting was performed using a rabbit polyclonal antibody specific to the phosphorylated form of Smad2. The same nitrocellulose membrane was reblotted and with an anti-STAT3

antibody to demonstrate equal protein loading. This data is representative of at least three different experiments.

**Figure 11:** Autophosphorylation of type II kinase in HaCat and GPI anchor mutated cells. HaCat or GPI M cells were left untreated (-) or treated (+) with 100 pM of TGF- $\beta$ 1 for 20 minutes. Precleared lysates were immunoprecipitated with 3  $\mu$ g/ml of anti-type II receptor antibody overnight. Following adsorption to protein A sepharose beads, 10  $\mu$ Ci of gamma  $^{32}$ P was added to the immunocomplexes and incubated for 30 minutes at 30°C to allow phosphorylation to occur. The reaction was halted by the addition of sample buffer and immunocomplexes were then subjected to SDS-PAGE under reducing conditions. This data is representative of two different experiments.

**Figure 12:** Schematic diagram representing the cloned sequence of the r150 protein.

**Figure 13:** HaCaT or 293 cells were transfected with r150 gene or the empty vector (pCMV sport 6) and cell lysates were fractionated by SDS-PAGE and transferred onto nitrocellulose membrane and immunoblotted with anti-CRD antibody. The Western blot shown is representative of four experiments (two each with HaCaT and 293 cells).

**Figure 14:** HaCaT or 293 cells were transfected with r150 gene or the empty vector (pCMV sport 6) or were left untransfected. Cells were allowed to recover for 24 hrs and were treated with 100 pM TGF- $\beta$ 1 for 30 minutes. Cell lysates were then Western blotted with an anti-phosphoSmad2 antibody.

**Figure 15:** HaCaT or 293 cells were transfected with r150 gene or the empty vector (pCMV sport 6) or were left untransfected. Cells were allowed to recover for 24 hrs and were treated with 100 pM TGF- $\beta$ 1 for 24 hours, or were left untreated. The luciferase activity was normalized to galactosidase activity obtained from a cotransfected CMV gal plasmid.

**Figure 16:** Schematic model of the mechanism by which r150 inhibits TGF- $\beta$  responses.

**Figure 17: Sequences of CD109, published by another, representing 150 putative variants. (Genbank accession number AF410459 and AAL84159.1.)**

**Figure 18: Alignment of  $\alpha_2$ -macroglobulin and r150 partial sequences.**

**EXAMPLE 1:**

**CHARACTERIZATION OF TGF- $\beta$ 1 BINDING PROTEIN DIFFERENT FROM OTHER TGF- $\beta$  RECEPTORS**

**METHODS**

**10 Cell Culture:**

Neonatal keratinocytes were prepared from foreskin tissue obtained at newborn male circumcision as described by Germain et al (1993). The keratinocytes were cultured in keratinocyte serum free medium (Gibco, Burlington, Ontario) and cells of third to fifth passage were used for experiments. The immortalized keratinocyte cell line, HaCaT, was obtained from Dr. Boukamp (Heidelberg, Germany), and the mink lung epithelial cells (Mv1Lu) were from ATCC. Both cell types were maintained in Dulbecco's Minimal Essential Medium (D-MEM) supplemented with 5% FBS, 1 mM sodium pyruvate, 2 mM glutamine, 100 U/ml penicillin, 100  $\mu$ g/ml streptomycin, and 0.25  $\mu$ g/ml amphotericin (Gibco, Burlington, Ontario). All cells were maintained at 37°C in an atmosphere of 5% CO<sub>2</sub>/air.

**Affinity labeling of cells:**

Iodination of TGF- $\beta$ 1 (Collaborative Biomedical) was done as described (Philip and O'Connor-McCourt, 1991). Affinity labeling technique was performed as detailed previously (Dumont et al, 1995). Briefly, cell monolayers were washed with ice-cold binding buffer (D-PBS or Dulbecco's phosphate-buffered saline with Ca<sup>2+</sup> and Mg<sup>2+</sup>, pH 7.4) containing 0.1% bovine serum albumin (BSA). Cells were incubated with 100- 200pM of <sup>125</sup>I-TGF- $\beta$ 1 for three hours at 4°C. In some experiments, incubations were done in the absence or presence of increasing

- 21 -

concentrations of unlabeled TGF- $\beta$  isoforms to determine the competition profiles of the receptors. The receptor-ligand complexes were cross-linked with 1mM Bis-(Sulfosuccinimidyl) suberate (BS<sup>3</sup>, Pierce). The reaction was stopped by the addition of glycine and the cells were solubilized, and separated on 3-11% polyacrylamid SDS gel. The results were analyzed by using autoradiography, followed by quantitative densitometry (Gel-Cypher, Lighttools Inc, Encinitas, CA or ImageQuant, Molecular Dynamics, Sunnyvale, CA)

*Temperature induced phase separation in Triton X-114 of r150 and hydrolysis by PIPLC:*

Temperature induced phase separation in Triton X-114 and PIPLC treatment was performed as described previously with modifications (Bordier, 1981). Keratinocytes were affinity labeled with 150 pM of <sup>125</sup>I-TGF- $\beta$ 1 and lysed in TBS (10 mM Tris-HCl, pH 7.5, 150 mM NaCl) containing 1% Triton X-114, 1 mM phenylmethylsulfonyl fluoride and protease inhibitor cocktail (200  $\mu$ g/ml BSA, 1  $\mu$ g/ml leupeptin, 10  $\mu$ g/ml benzamide, 10  $\mu$ g/ml soyabean trypsin inhibitor, and 2  $\mu$ g/ml pepstatin) for 60 minutes at 4°C. The cell lysates were centrifuged at 13 000 x g for 15 minutes at 4°C. The Triton X-114 soluble material was incubated at 30°C for 10 minutes followed by a 10 minute centrifugation at 13 000 x g at room temperature to separate the detergent rich phase from the aqueous detergent poor phase. An aliquot (20%) from each phase was precipitated with ethanol/acetone, and analyzed by SDS-PAGE and autoradiography. The remaining 80% of the detergent phase was utilized to determine the effect of PIPLC on the detergent solubility of r150 using the method of Lisanti et al (1988). Briefly, the GPI-anchored protein enriched detergent phase was incubated with or without 0.6 U/ml of PIPLC (Roche Diagnostics) for one hour at 37°C with mild agitation. Temperature induced phase separation was then repeated. Both the aqueous and detergent phases were precipitated by adding ethanol/acetone, and subjected to SDS-PAGE and autoradiography.

*Affinity labeling of soluble r150:*

Neonatal keratinocytes were harvested by treating confluent monolayers with Hanks' balanced salt solution containing 5 mM EDTA (pH 7.5). The

cell pellet was washed with D-PBS and treated with 0.6 U/ml of PIPLC or left untreated, for one hour at 37°C with mild agitation.

The supernatant containing the released GPI-anchored proteins was collected and concentrated by Centricon 30 (Amicon). Aliquots of the concentrated supernatant were affinity labeled with 150 pM of  $^{125}\text{I}$ -TGF- $\beta$ 1 in the absence or presence of excess unlabeled TGF- $\beta$  (7.5 nM) and analyzed by SDS-PAGE as described above except that the solubilization step was omitted.

*$^{125}\text{I}$ -TGF- $\beta$ 1 binding to Mv1Lu cells*

To test whether soluble r150 regulates the binding of TGF- $\beta$  to its receptors, the supernatant obtained from PIPLC treated HaCaT cells were used in a  $^{125}\text{I}$ -TGF- $\beta$ 1 Mv1Lu binding assay. The HaCaT cells, which display the r150 with identical properties as the neonatal keratinocytes (Tam et al, 1998), were grown in T-25cm<sup>2</sup> tissue culture flasks (Falcon) were left untreated or treated with PIPLC, and the resulting supernatants were concentrated by Centricon 30 (Amicon). Mv1Lu cells were incubated with 50 pM of  $^{125}\text{I}$ -TGF- $\beta$ 1 in the absence or presence of increasing doses of the concentrated supernatant for three hours at 4°C. The cells were washed, solubilized and the bound radioactivity was determined by a gamma counter.

To rule out the possibility that any alteration in  $^{125}\text{I}$ -TGF- $\beta$ 1 binding caused by the supernatants was not due to the presence of TGF- $\beta$ 1, the above binding assay was also done in the presence of the supernatant treated with an anti-TGF- $\beta$ 1 antibody. The supernatant was incubated with the antibody (15  $\mu\text{g}/\text{ml}$ ) overnight at 4°C. It was then precleared of immune complexes and excess antibody that may interfere with the assay, by incubating with a protein A Sepharose slurry (PharmaciaBiotech) for two hours before addition to the assay. The TGF- $\beta$ 1 antiserum (obtained from Dr.M. O'Connor-McCourt ; Moulin et al, 1997) was purified on a HiTrap Protein G column (PharmaciaBiotech) following standard procedures. The specificity of the antibody was verified by Western Blot analysis

*Determination of TGF- $\beta$  concentration in HaCaT supernatant:*



- 23 -

The concentration of TGF- $\beta$ 1 was quantitated in the supernatants from HaCaT cells left untreated or treated with PIPLC using a mink lung epithelial cells-luciferase assay as described by Nunes et al (1996). This quantitative bioassay for TGF- $\beta$  is based on the ability of TGF- $\beta$  to induce the expression of plasminogen activator inhibitor type 1 (PAI-1) gene. The mink lung epithelial cells stably transfected with an expression construct containing a truncated PAI-1 promoter fused to the luciferase reporter gene were provided by Dr. D.B. Rifkin (New York University Medical Center). Briefly, these cells were incubated with varying doses of HaCaT supernatant, and the luciferase activity (expressed as relative light units) was quantitated by a Berthold Luminometer. Recombinant TGF- $\beta$ 1 (Austral Biochemicals; 0.5 pM-50 pM) was used to create a standard curve.

*Immunoaffinity Chromatography and Immunoblotting of r150:*

Neonatal keratinocytes were harvested by treating confluent monolayers with Hanks' balanced salt solution containing 5 mM EDTA (pH 7.5). The cell pellet was washed with D-PBS and treated with 0.6 U/ml of PIPLC for one hour at 37°C with mild agitation. The supernatant containing the released GPI-anchored proteins was purified through a TGF- $\beta$ 1 affinity column (made available by Dr. M. O'Connor-McCourt, Montreal, Quebec). The column was prepared by incubating one milligram of TGF- $\beta$ 1 in a 200 mM HCO<sub>3</sub> (pH 8.3) buffer containing 30% (v/v) of n-propanol with five milligrams of Reacti-gel (Pierce) for 72 hours at 4°C. The reaction was stopped by the addition of 200  $\mu$ l of 2 M TBS (pH 7.4), and the gel was washed to remove any unbound TGF- $\beta$ 1. The supernatant containing the GPI-anchored medium was loaded on the column equilibrated with 50 mM Tris (pH 7.4) and eluted with a 10 mM citrate/300 mM NaCl buffer (pH 2.5). Fractions of 0.5 ml were collected, and each fraction was analyzed for binding to TGF- $\beta$ 1, and immunoblotted for the presence of GPI anchor.

To verify binding to TGF- $\beta$ 1, an aliquot from each fraction was affinity labeled with 150 pM of <sup>125</sup>I-TGF- $\beta$ 1 and analyzed by SDS-PAGE and autoradiography as described above, except that the solubilization step was omitted.

- 24 -

The fraction containing r150 and adjacent fractions were immunoblotted with the anti-CRD antibody to detect the GPI anchor. The anti-CRD antibody is specific to the inositol 1,2 cyclic monophosphate moiety, known as the "cross-reacting determinant" (CRD) which is exposed in GPI-anchored proteins that have been hydrolyzed by PIPLC. The antibody obtained from Oxford GlycoSystems (Wakefield, MA), was raised against the inositol 1,2 cyclic monophosphate moiety of the trypanosome variant surface glycoprotein (VSG). Samples were analyzed on 3-11% polyacrylamide gradient SDS gels and transferred to nitrocellulose membrane. The membrane was blocked in TBS-T (30 mM Tris, 150 mM NaCl, pH 7.5, 0.5% Tween 20) containing 5% non-fat dry milk and was incubated overnight with 4 µg/ml of the anti-CRD antibody at 4°C. The blots were washed in TBS-T and incubated for three hours at room temperature with the alkaline phosphatase conjugated secondary antibody (1:1500) (Roche Diagnostics). The membrane was then subjected to chemiluminescence analysis (CDP-Star) as detailed by the manufacturers (Roche Diagnostics).

#### **Immunoprecipitation of r150:**

Two different anti-CRD antibodies (i) Oxford GlycoSystems antibody and (ii) an antibody raised against a mammalian GPI-anchored pig membrane dipeptidase (MDP) were used for immunoprecipitation studies. The latter antibody was isolated from the bulk of the anti-MDP antiserum by fractionation on a column of the immobilized form of the trypanosome variant surface glycoprotein. Both anti-CRD antibodies are specific to the inositol 1,2-cyclic monophosphate and have been well characterized (Zamze et al, 1988; Broomfield and Hooper, 1993). Cells were affinity labeled with 200 pM <sup>125</sup>I-TGF-β1 and cell extracts were incubated with the anti-CRD antibody. The resulting immune complexes were treated with protein A Sepharose (Pharmacia-Biotech) slurry and the beads were pelleted by centrifugation, and were analyzed by SDS-PAGE under reducing conditions followed by autoradiography.

## **RESULTS**

**Binding Affinity of r150 for TGF- $\beta$  isoforms:**

The present inventors have previously reported that in addition to the types I, II and III receptors, keratinocytes express a novel GPI-anchored TGF- $\beta$ 1 binding protein r150 which forms a heteromeric complex with the TGF- $\beta$  signaling receptors (Tam et al, 1998). Since this protein has the potential to regulate TGF- $\beta$  signaling, it was further characterized. Here was determined the relative affinity of r150 for the three TGF- $\beta$  isoforms and, this affinity for TGF- $\beta$ 1 approximates that of the TGF- $\beta$  signaling receptors, which suggests that r150 is predominantly a TGF- $\beta$ 1 binding protein. Keratinocytes affinity labeled with  $^{125}$ I-TGF- $\beta$ 1 in the absence or presence of increasing concentrations of unlabeled TGF- $\beta$ 1, - $\beta$ 2 or - $\beta$ 3, were analyzed by SDS-PAGE and autoradiography (Figure 1a), and competition curves were created from the autoradiogram for r150, type I and type II receptors using quantitative densitometry (Figures 1b, c, and d). r150 is not sensitive to reducing agents since its migration pattern is identical when the SDS-PAGE analysis is done under non-reducing (Figure 1a) or reducing conditions (Tam et al, 1998). The half-maximal inhibition of  $^{125}$ I-TGF- $\beta$ 1 binding was determined from the competition curves as the TGF- $\beta$  isoform concentration at which the inhibition was 50% of that observed when no unlabeled ligand was present (Table 1). The concentration of unlabeled TGF- $\beta$ 1 required for half maximal inhibition of  $^{125}$ I-TGF- $\beta$ 1 binding by r150 is only 1.2 and 1.3 times higher than that required by type I and II receptors respectively. Although r150 also binds TGF- $\beta$ 3, it does so with a much lower affinity as compared to the types I and II receptors since it requires a six-fold higher concentration of TGF- $\beta$ 3 to reach half-maximal inhibition of  $^{125}$ I-TGF- $\beta$ 1 binding than the types I or II receptors. Unlabeled TGF- $\beta$ 2, even at 40 times excess concentrations minimally inhibited  $^{125}$ I-TGF- $\beta$ 1 binding of r150.

***Partitioning of the membrane bound and released r150 in Triton X-114:***

In order to ascertain that the membrane bound r150 is hydrophobic as expected of a GPI-anchored protein and that the released r150 behaves as a hydrophilic soluble protein, the temperature dependent phase separation property of the non-ionic detergent Triton X-114 was used. Phase separation using Triton-X 114 results in the partitioning of hydrophilic proteins into

the aqueous detergent poor phase while integral membrane proteins and lipid attached proteins partition into the detergent rich phase. This procedure has been useful in distinguishing between the amphipathic (membrane bound) and hydrophilic (released from cell surface) forms of GPI-anchored proteins (Hooper, 1992).

5 Affinity labeled keratinocytes were subjected to Triton X-114 partitioning and the detergent rich phase containing hydrophobic proteins and the detergent poor phase containing the hydrophilic proteins were analyzed by SDS-PAGE. As expected of a GPI-anchored protein, r150 partitioned predominantly into the detergent rich phase, along with the transmembrane type I, II and III receptors  
10 (Figure 2A). The partitioning of soluble r150 in Triton X-114 was then tested. When the detergent rich phase containing the membrane bound affinity labeled r150 was left untreated or treated with PIPLC, and the temperature-induced phase separation was repeated, the aqueous phase of the sample treated with PIPLC was enriched in r150 while that of the sample left untreated contained only low amounts of r150  
15 (Figure 2B). These results strongly indicate that the PIPLC-released r150 is indeed hydrophilic. In contrast, the detergent phase of samples left untreated with PIPLC contained the major portion of r150 while the detergent phase of samples treated with PIPLC contained minimal amounts of r150 (data not shown).

#### **The soluble r150 binds TGF- $\beta$ 1**

20 Was next examined whether r150 released from the cell surface is capable of binding TGF- $\beta$ 1.

Data shown in Figure 3a demonstrate that soluble r150 in the supernatant obtained from keratinocytes treated with PIPLC could be affinity labeled with  $^{125}$ I-TGF- $\beta$ 1. This binding was specific since it was markedly reduced when the  
25 labeling was done in the presence of unlabeled TGF- $\beta$ 1. unlabeled TGF- $\beta$ 1 did not exhibit any competition for these complexes.

The low molecular weight bands below 97.4 kDa appear to be nonspecific since unlabeled TGF- $\beta$ 1 did not exhibit competition for these complexes in a reproducible manner. The fact that released r150 binds TGF- $\beta$ 1 indicates that

r150 is capable of binding the ligand in the absence of type I, II and III TGF- $\beta$  receptors or an intact membrane structure. Interestingly, detectable amounts of r150 were observed in the supernatant not treated with PIPLC, which led us to suspect that there might be an endogenous phospholipase capable of releasing r150.

5                   That the soluble r150 can inhibit TGF- $\beta$ 1 binding to TGF- $\beta$  receptors was demonstrated using a binding assay. As seen in Figure 3b, the supernatant from PIPLC treated keratinocytes competed in a dose dependent fashion for  $^{125}$ I-TGF- $\beta$ 1 as seen by decreased binding to MvLu1 cells. The supernatant from a T-25cm<sup>2</sup> flask treated with PIPLC inhibited binding by 33%  
10   (p<0.005) and 50% (p<0.04) at doses of 1 and 2 respectively (approximately  $1 \times 10^6$  cells, represented as an arbitrary unit of "1" in Figure 3b). The inhibition of binding with the supernatant from cells not treated with PIPLC is consistent with the observation that detectable amounts of r150 is present in this supernatant, alluding to the presence of an endogenous phospholipase capable of releasing r150 (Figure  
15   3a; also see below, Figure 5). This inhibition of  $^{125}$ I-TGF- $\beta$ 1 binding to MvLu1 cells corresponded to 15%, and 31% (p<0.03 in both cases), respectively for doses 1 and 2. In the PIPLC treated supernatants, the inhibition of binding at doses 1 and 2 was significantly higher (p<0.03 in both cases) than in the untreated supernatants.

                  To rule out the possibility that the competition observed by  
20   PIPLC treated supernatant was due to TGF- $\beta$ , the supernatant was neutralized with anti-TGF- $\beta$ 1 antibody prior to being used in the binding assay. Neutralization with this antibody had no effect on the inhibition by the r150 enriched supernatant (Figure 3c). In contrast, 100 pM TGF- $\beta$ 1 markedly inhibited  $^{125}$ I-TGF- $\beta$ 1 binding and this binding could be neutralized by anti-TGF- $\beta$ 1 antibody but not by non-immune IgG.  
25   Furthermore, using a PAI-luciferase assay no TGF- $\beta$  was detected in the supernatants of cells untreated or treated with PIPLC (data not shown). Taken together, these results suggest that the released form of r150 is capable of binding to TGF- $\beta$ 1 and modulating ligand binding to TGF- $\beta$  receptors.

*Identification of a GPI-anchor in r150:*

Although r150 is sensitive to PIPLC, it is possible that it is not itself GPI-anchored, but is associated with a protein that is GPI anchored. Also, it is conceivable that it is a complex of two lower molecular weight proteins which became inadvertently cross-linked during the affinity labeling procedure. In order to eliminate these possibilities, Western blot analysis of r150 was performed after its release from the cell membrane using an anti-CRD antibody specific for an epitope which becomes exposed in GPI anchored proteins only upon treatment with PIPLC.

Keratinocytes were treated with PIPLC and the supernatant was purified on a TGF- $\beta$ 1 affinity column. Analysis of fractions by affinity labeling and SDS-PAGE demonstrated that the fraction 21, but not adjacent fraction (represented by fraction 18) contained a 150 kDa TGF- $\beta$ 1 binding protein (Figure 4a). The binding of  $^{125}$ I-TGF- $\beta$ 1 to this protein is specific since it was blocked in the presence of 5 nM TGF- $\beta$ 1. These results confirm that soluble r150 binds TGF- $\beta$ 1.

When the fractions were analyzed by Western blotting with the anti-CRD antibody, it was revealed that fraction 21, but not other fractions contained a protein of relative molecular weight of 150 kDa which was recognized by the anti-CRD antibody. Detection of a 150 kDa protein by the anti-CRD antibody in the absence of chemical cross-linking demonstrates that r150, but not an associated protein, contains a GPI-anchor and, that r150 does not represent two smaller proteins which got inadvertently cross-linked (Figure 4b). Figure 4b also shows that r150 was not detectable in an adjacent fraction (fraction 18).

**Evidence to indicate that an endogenous phospholipase C releases r150 in human keratinocytes:**

Next, whether the anti-CRD antibody can immunoprecipitate r150 and/or coprecipitate the types I and II TGF- $\beta$  receptors was tested. During the course of these studies, it was observed that the anti-CRD antibody immunoprecipitated r150 from  $^{125}$ I-TGF- $\beta$ 1 labeled keratinocytes, even in the absence of PIPLC treatment (Figure 5). This result was reproducible using two different anti-CRD antibodies: the Oxford GlycoSystems antibody that is raised against the CRD epitope of variant surface glycoprotein of *Trypanosoma brucei*

- 29 -

(Oxford GlycoSystems), and the antibody specific for the CRD epitope of the porcine membrane dipeptidase (MDP, Broomfield and Hooper, 1993). Since both anti-CRD antibodies are specific for the inositol 1,2-cyclic monophosphate epitope exposed only upon PIPLC treatment, recognition by the two antibodies in the absence of PIPLC treatment indicates that an endogenous phospholipase C cleaved r150 to expose this epitope. The anti-CRD antibody not only precipitated the r150, but also coprecipitated the types I and II receptors. Interestingly, the intensities of the types I and II bands were much stronger than that of the r150 itself. The coimmunoprecipitation of the types I and II receptors demonstrates the heteromeric complex formation of r150 with those receptors (Figures 5a, b) which confirms the inventors' previous finding (Tam et al, 1998). The immunoprecipitation with the anti-CRD antibody is specific because the precipitation of labeled complexes is efficiently blocked when the PIPLC hydrolyzed form of MDP which contains the epitope to which the antibody was raised against was included in the reaction (Figure 5b). These complexes are not detected when the cells were affinity labeled with <sup>125</sup>I-TGF-β2 because r150 has a much lower affinity for TGF-β2 than for the TGF-β1 and TGF-β3 isoforms in these cells (Figure 5c).

## DISCUSSION

The present inventors have shown that a novel 150 kDa TGF-β1 accessory receptor (r150) forms a heteromeric complex with the TGF-β signaling receptors on human keratinocytes (Tam et al. 1998). This accessory receptor was described as GPI-anchored based on its sensitivity to PIPLC. Here it is demonstrated that the GPI-anchor is contained in r150 itself and not on an associated protein and that it binds TGF-β1 with an affinity similar to those of the types I and II TGF-β receptors. In addition, evidence is provided that r150 is released from the cell surface by an endogenous phospholipase C. The most important finding in the present work is that the released (soluble) form of r150 binds TGF-β1 independent of the signaling receptors.

r150 has been characterized as GPI-anchored, based on its sensitivity to phosphatidylinositol specific phospholipase C (PIPLC). In order to prove that the GPI-anchor is present in the r150 itself, it was necessary to rule out other

possibilities, namely: (i) r150 is not itself GPI-anchored, but is tightly associated with a protein that is GPI-anchored. Upon PIPLC treatment, the associated GPI-anchored protein is cleaved which results in the release of both the proteins into the supernatant. This has been shown to be the case for lipoprotein lipase which was initially identified as GPI-anchored protein based on its sensitivity to PIPLC; but it was later found that its PIPLC sensitivity was a result of close association with a GPI-linked heparan sulfate proteoglycan (Chajek-Shaul et al, 1989). (ii) r150 is a noncovalently associated complex of two lower molecular weight proteins whose combined molecular weights equate 150 kDa, of which one component is GPI-anchored. During affinity labeling, the two proteins get inadvertently crosslinked by the chemical crosslinker BS<sup>3</sup>, and thus upon analysis by SDS-PAGE, the cross-linked complex is detected at 150 kDa.

By immunoblotting the purified, soluble form of the r150 with the anti-CRD antibody that can specifically recognize the epitope exposed by the cleavage of the GPI-anchor by PIPLC, the above two possibilities were eliminated. Elution from a TGF- $\beta$ 1 affinity column and detection as a 150 kDa protein in the absence of cross-linking, together with its ability to be recognized by the anti-CRD antibody, prove that r150 has a relative molecular weight of 150 kDa and that the GPI-anchor is contained in r150 itself. There are two GPI-anchored proteins expressed in mammalian tissues that have similar molecular weights as r150. These include an isoform of NCAM (140 kDa) (Rosen et al, 1992) and ceruloplasmin (135 kDa) (Patel and David, 1997). However, immunoprecipitation with antibodies specific to these proteins did not immunoprecipitate r150 affinity labeled with <sup>125</sup>I-TGF- $\beta$ 1 (data not shown).

The soluble r150 is capable of binding to TGF- $\beta$ 1 as shown by affinity labeling of the released protein, and retention on the TGF- $\beta$ 1 affinity column. This suggests that r150 can bind TGF- $\beta$  in the absence of types I, II and III receptors or an intact membrane. That soluble r150 has the potential to modulate TGF- $\beta$  binding to its receptors was demonstrated by its ability to inhibit <sup>125</sup>I-TGF- $\beta$ 1 binding to receptors on mink lung cells. Although studies of the inhibition of TGF- $\beta$  binding to its receptors by r150 used cellular supernatant and not purified r150, this inhibition



is most likely due to r150 itself, since there was no measurable TGF- $\beta$  and neutralizing anti-TGF- $\beta$  antibody had no effect on this inhibition. The inhibition obtained by the supernatant not treated by PIPLC is likely due to endogenous release of r150 (Figure 5). As expected, exogenous addition of PIPLC resulted in significantly higher inhibition of binding since more r150 will be released. In addition, r150 is the major TGF- $\beta$  binder released by PIPLC (Figure 2). The other potential binder,  $\alpha$ 2-macroglobulin is unlikely to be released in sufficient quantity during the one hour incubation.

That soluble r150 binds TGF- $\beta$ 1 is reminiscent of what is observed of the ectodomain of type III receptor which has been shown to be released by a not yet characterized mechanism (Lopez-Casillas et al, 1994; Philip et al, 1999). The soluble r150 may act as an antagonist by preventing the binding of TGF- $\beta$  to the signaling receptors as has been suggested for the soluble type III receptor (Lopez-Casillas et al, 1994). This is supported by our finding which suggests that the soluble r150 inhibits  $^{125}$ I-TGF- $\beta$ 1 binding to receptors on mink lung cells. But unlike the type III receptor, r150 would antagonize TGF- $\beta$ 1 activity in an isoform specific manner, since it has a low affinity for TGF- $\beta$ 2 and a moderate affinity for TGF- $\beta$ 3, as determined by competition experiments using unlabeled TGF- $\beta$  isoforms. Furthermore, affinity cross-link labeling of keratinocytes with  $^{125}$ I-TGF- $\beta$ 2 or  $^{125}$ I-TGF- $\beta$ 3 did not demonstrate labeling of r150 (data not shown). Recently, a soluble type I receptor has been cloned from a rat kidney cDNA library (Choi, 1999). In contrast to the soluble type III receptor, the soluble type I receptor requires the co-expression of the type II receptor in order to bind TGF- $\beta$ . However, it appeared to potentiate TGF- $\beta$  signal transduction and the author has suggested that this potentiation may be due to the stabilization of the heteromeric TGF- $\beta$  signaling receptor complex. The observation that the soluble r150 can bind the ligand independently of signaling receptors, and modulate TGF- $\beta$  binding to its receptors (present work), together with the fact that the membrane bound r150 binds TGF- $\beta$ 1 and forms heteromeric complex with the type I and II receptors (Tam et al, 1998) raise the possibility that r150 in its membrane bound or soluble form may act as antagonist of TGF- $\beta$  signaling by regulating ligand availability, or stability of the signaling receptor complex or by directly affecting the signal transduction process.

It is predicted that r150 has dual roles in TGF- $\beta$  signaling depending on whether it exists as a cell surface anchored protein or as the soluble form. Consequently, a potential mechanism for the regulation of its action would be the hydrolysis of the GPI-anchor. Both the release of the protein from the cell surface and the ability of the soluble form to sequester TGF- $\beta$  may modulate TGF- $\beta$  receptor function. Such a possibility *in vivo* is supported by our observation that an endogenous phospholipase C releases r150 from the cell surface. Since the anti-CRD antibody only recognizes GPI-proteins released from the cell surface by PIPLC, and the cells were not pretreated with PIPLC, the results indicate that there is an endogenous phospholipase C in keratinocytes capable of hydrolyzing r150 at the same site as the PIPLC. The soluble r150 identified in the cellular extract is not due to protease activity since proteolytic cleavage will result in a protein of lower molecular weight. Neither can it be due to failure of cells to add the GPI anchor during synthesis, since the antibody will not recognize that protein. It is of interest to note here that r150 was detectable upon overexposure of films in the PIPLC untreated aqueous fractions obtained by Triton X-114 partitioning (data not shown).

Although the presence of a mammalian PIPLC has not been definitively established, the activity of PIPLC-like enzymes have been implicated in the insulin signaling pathway in rat liver (Satiel, 1996). Furthermore, Movahedi and Hooper (1997) have demonstrated that the insulin stimulated release of GPI-anchored proteins from differentiated 3T3-L1 adipocytes occurs via the action of an endogenous phospholipase C. We choose the general term of phospholipase C for the enzyme that releases the r150 in keratinocytes, since the identity and specificity of the enzyme is not confirmed. Nevertheless, our results suggest that the release of r150 involves an enzyme that hydrolyzes r150 at the same site as PIPLC. PIPLD hydrolyze the GPI-anchor at a different site from PIPLC which does not result in the formation of the inositol 1,2-cyclic monophosphate, and therefore the anti-CRD antibodies that we used in this study cannot recognize PIPLD cleaved proteins (Broomfield and Hooper, 1993). However, PIPLD is expressed in mammalian serum and has been shown to be present in keratinocytes (Xie et al, 1993). Lin et al. (2002), who describe a cell surface antigen named CD109, which appears to be very similar to r150, propose that some GPI anchors are acylated on inositol, because of CD109

- 33 -

sensitivity to phospholipidase D. It is possible that the activity of both enzymes may be involved in regulating the cell surface expression of r150 on human keratinocytes.

GPI-anchored proteins have been reported to bind TGF- $\beta$  on certain cell lines (Cheifetz and Massague, 1991). More recently, the present inventors reported the presence of GPI-anchored TGF- $\beta$  binding proteins on early passage human endometrial stromal cells (Dumont et al, 1995) and human skin fibroblasts (Tam and Philip, 1998). Both endometrial stromal cells and skin fibroblasts displayed a 180 kDa GPI-anchored TGF- $\beta$ 1 binding protein and a 65 kDa TGF- $\beta$ 2 binding protein. However, whether the GPI anchor is present in the proteins themselves has not been ascertained for any of them, and the identities of these proteins remain unknown. Interestingly, GPI-anchored proteins have been implicated in the maintenance of the epidermis. When the expression of GPI-anchored proteins was abrogated in the skin by the tissue specific deletion of the *PIG-A* gene, a gene essential for GPI-protein synthesis, mutants died shortly after birth and their skin was wrinkled and scaly in comparison to that of the wild type (Tarutani et al, 1997). Since TGF- $\beta$  has an important role in epidermal homeostasis, it is conceivable that the GPI-function which is compromised in these mutants is related to dysregulated TGF- $\beta$  action due to the loss of r150.

Our results from the competition experiments demonstrating that r150 has high affinity for the TGF- $\beta$ 1 isoform suggest that it is an endogenous ligand for this protein. Whether the membrane bound or soluble form of r150 acts as scavenger receptors regulating ligand availability, whether they participate directly in the modulation of downstream signaling, or if the release of the soluble form is a regulated event *in vivo*, remain to be determined. Identification of its structure should facilitate resolution of these issues. Elucidating the mechanism by which r150 functions as an accessory molecule in TGF- $\beta$  signaling in keratinocytes may be critical to understanding the molecular mechanisms underlying the regulation of TGF- $\beta$  action.

### Example 2

## CREATION OF KERATYNOCYTE CELL LINE MUTANTS DEFICIENT IN GPI ANCHOR BIOSYNTHESIS

### MATERIALS AND METHODS

- 5 **Cell Culture:** An immortalized human keratinocyte cell line, HaCat, was obtained from Dr. P. Boukamp (Hedidelberg, Germany). HaCat is a spontaneously immortalized cell line which displays no major differences in differentiation as compared to normal keratinocytes. It possesses a transformed phenotype, but is not tumourigenic (Boukamp et al, 1988). HaCat cells were cultured in D-MEM containing 10% fetal  
10 bovine serum, 1 mM sodium pyruvate, 2 mM glutamine and 100 U/ml penicillin, 100 µg/ml streptomycin, and 0.25 µg/ml amphotericin (all Gibco, Burlington, Ontario).

*Doubling time:*  $8 \times 10^5$  cells were seeded at in 60 mm dishes (Falcon) in duplicate. At indicated times, cells were trypsinized and counted using a hemocytometer

- Affinity labeling of cells:** Affinity cross-link labeling techniques were performed as  
15 detailed by Dumont et al (1995). Briefly, monolayers of cells were washed with ice cold binding buffer [PBS with  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$ , pH 7.4 (D-PBS) containing 0.1% BSA] three times over a thirty minute period. Cells were then incubated at 4°C for three hours with 100 pM of  $^{125}\text{I}$ -TGF- $\beta$ 1 in the presence or absence of excess non-radioactive TGF- $\beta$ 1 (Austral Biochemical, Genzyme Inc. or R & D Systems  
20 respectively). The receptor-ligand complexes were then cross-linked with 1mM Bis(Sulfosuccinimidyl)suberate (BS3) (Pierce, Rockford IL). After 10 minutes, the reaction was stopped by the addition of 500 mM glycine and further incubated for 5 minutes. The cells were then washed twice with D-PBS and lysed with solubilization  
25 buffer (20 mM Tris-HCl, pH 7.4 containing 1% Triton X-100, 10% glycerol, 1 mM EDTA, 10 µM phenylmethylsulfonyl fluoride (PMSF), 200 µg/ml BSA, 1 µg/mL leupeptin, 10 µg/ml soyabean trypsin inhibitor, 10 µg/ml benzamide and 2 µg/ml pepstatin). The solubilized material was collected and 1/5 volume of 5X electrophoresis sample buffer (0.25 M Tris-HCl, pH 6.8, 5% SDS, 50% glycerol and trace bromophenol blue) was added. The samples were run on a 1.5 mm-thick 3%-

11% SDS-PAGE under nonreducing or reducing (in the presence of 5%  $\beta$ -mercapto ethanol) conditions. Results were analyzed using autoradiography followed by quantitative densitometry. [ $^{14}\text{C}$ ] methylated molecular weight protein markers included myosin (H-chain) (200-220 kDa), phosphorylase-b (97.4 kDa), bovine serum albumin (68 kDa), ovalbumin (43 kDa), and carbonic anhydrase (29 kDa),  $\beta$ -lactoglobulin (18.4 kDa) and lysozyme (14.3 kDa) (Gibco or Amersham-Pharmacia Biotech).

**Affinity labeling of soluble r150.** Confluent monolayers of keratinocytes were treated with 0.6 U/ml of PIPLC for one hour at 37°C with mild agitation. The supernatant containing the released GPI-anchored proteins was collected and concentrated by Centricon 30 (Amicon). To perform affinity labeling of the soluble form of r150, aliquots of the concentrated supernatant were affinity cross-link labeled with 100-150 pM of  $^{125}\text{I}$ -TGF- $\beta$ 1 in the absence or presence of excess unlabeled TGF- $\beta$ 1 as described above except that the solubilization step was omitted. Binding complexes were analyzed by SDS-PAGE and autoradiography.

**Isolation and cloning of a keratinocyte cell line mutated in GPI anchor biosynthesis.** The preparation of a keratinocyte cell line mutated in GPI anchor biosynthesis was performed as described by Stevens (1999). HaCat cells grown to 50-60% confluence in a T-75cm<sup>2</sup> flask were treated with 300  $\mu\text{g}/\text{ml}$  of ethylmethane sulfonate (EMS) for 24 hours at 37°C in an atmosphere of 5%  $\text{CO}_2/\text{air}$ . EMS is one of the most common chemical mutagens used to generate cells that are defective in GPI anchor biosynthesis (Sega, 1984). Cells were then allowed to recover for an additional 48 hrs. in normal growth medium. Fluorescence activated cell sorting (FACS) was performed to select those cells which lost the expression of GPI-anchored proteins from their cell surface. Since the identity of the r150 was not known, GPI anchor deficient cells were negatively sorted for another GPI-anchored protein, CD59. Briefly, cells were trypsinized and approximately  $2 \times 10^6$  cells were resuspended in one milliliter of FACS buffer (PBS containing 1% FBS) and incubated with 4  $\mu\text{g}$  of fluorescein isothiocyanate (FITC)-conjugated mouse anti-human CD59 monoclonal antibody (BD Pharmagins, Mississauga, ON.) or mouse FITC-conjugated IgG<sub>2a</sub> monoclonal antibody for 45 minutes on ice. After the incubation, the cell pellet

was washed once and resuspended in one milliliter of FACS buffer. Propidium iodide (3 µg/ml) (Sigma) was added to the cells prior to sorting which was performed using the FACS Vantage (Becton Dickinson). Cell sorting was repeated two additional times. At the third and final FACS, a single CD59 negative cell was seeded into each well of a 96 well plate using the FACS Vantage automated cell deposition unit. In the surviving clones, the cell surface expression of CD59 was analyzed by flow cytometry to determine which cells expressed deficient levels of GPI-anchored proteins. Affinity cross-link labeling with <sup>125</sup>I-TGF-β1 was performed to characterize the expression of the r150.

- 10 **Luciferase assay.** The p3TP-Lux and CMVβ-galactosidase (CMVβ-gal) gene constructs were gifts from Dr. O'Connor-McCourt (Biotechnology Research Institute, Montreal, Quebec). The p3TP-Lux construct contains three tetradecanoyl phorbol acetate (TPA)-responsive elements and TGF-β responsive elements from the PAI-1 promoter fused to the luciferase reporter gene (Wrana et al, 1992). Briefly, 2.5 x 10<sup>6</sup>
- 15 HaCat cells, plated in a 12 well plate, were transiently transfected with 1 µg each of p3TP-Lux and CMVβ-gal cDNAs using the Superfect reagent as recommended by the manufacturers (Qiagen, Mississauga, Ontario). The cells were allowed to recover overnight and were serum starved for three to four hours prior to treatment with various doses of TGF-β1 or TGF-β2 for the indicated times (Austral Biochemicals).
- 20 For the luciferase assay, cells were solubilized in 150 µl of lysis buffer (BD Pharmagin, Mississauga, Ontario) for 30 minutes at 4°C. In an opaque 96 well plate, 45 µl of lysate was added to 10 µl of ATP cocktail [0.1 M ATP, 0.5 M KH<sub>2</sub>PO<sub>4</sub> (pH 7.8), 1 M MgCl<sub>2</sub>]. Luciferase activity was measured upon addition of 100 µl of 25 mM luciferin (Roche Diagnostics) in 0.1 M KH<sub>2</sub>PO<sub>4</sub> (pH 7.8) using an EG & G Berthold
- 25 Microplate Luminometer. For the β-galactosidase assay, in a 96 well plate, 5 µl of the lysate was added to 100 µl of β-gal buffer (60 mM Na<sub>2</sub>HPO<sub>4</sub>, 40 mM NaH<sub>2</sub>PO<sub>4</sub>, 10 mM KCl, 1 M MgCl<sub>2</sub> and 50 mM β-mercaptoethanol) containing 1.5 ng/ml of ortho-nitrophenyl β D-galactopyranoside (ONPG) (Sigma). The samples were incubated at 37°C until a satisfactory colour reaction was obtained. The colour reaction was
- 30 then measured at 420 nm using a spectrophotometric plate reader (Molecular Dynamics, Sunnyvale, CA). Values were derived from transfections performed in duplicate or triplicate and all experiments were performed at least three times.

- 37 -

**In vitro kinase assay.** HaCat and GPI anchor mutated cells grown in 75cm<sup>2</sup> flasks were serum starved overnight and left untreated or treated with 100 pM TGF- $\beta$ 1 for 20 minutes at 37°C. The cells were scraped in 200  $\mu$ l of lysis buffer (50 mM Tris pH 7.5, 150 mM NaCl, 50mM NaF, 50 mM  $\beta$ -glycerophosphate, 1 mM sodium orthovanadate, 1 mM DTT, 5 mM EDTA pH 8.0, 1 % NP-40, 10 % glycerol, 10  $\mu$ M PMSF, 200  $\mu$ g/ml BSA, 1  $\mu$ g/mL leupeptin, 10  $\mu$ g/ml soyabean trypsin inhibitor, 10  $\mu$ g/ml benzamide and 2  $\mu$ g/ml pepstatin) and further incubated in the lysis buffer for 30 minutes on ice. The cells were pelleted and the solubilized extract was precleared with 40  $\mu$ l of protein A Sepharose slurry (Amersham-Pharmacia-Biotech) and 3  $\mu$ g/ml of polyclonal rabbit IgG antibody for two hours at 4°C. One mg of protein was incubated with 3  $\mu$ g/ml of anti -type II TGF- $\beta$  receptor antibody (Santa Cruz) overnight at 4°C. The immune complexes were then incubated with 50  $\mu$ l of protein A Sepharose slurry for two hours at 4°C. The beads were pelleted and washed three times with 0.5 ml of lysis buffer and one time with 0.5 ml of kinase buffer [50 mM Tris (pH 7.4), 10 mM MgCl<sub>2</sub>, 1 mM CaCl<sub>2</sub>]. After the last wash, 20  $\mu$ l of kinase buffer was added to the pellets and incubated with 10  $\mu$ Ci of gamma<sup>32</sup>P-ATP (NEN) for 30 minutes at 30°C. The reaction was halted by the addition of 5  $\mu$ l of 5X sample buffer (0.25M Tris-HCl, pH 6.8, 5% SDS, 50% glycerol and trace bromophenol blue) containing 5%  $\beta$ -mercaptoethanol and boiled for 5 minutes. The extracts were separated on a 7.5% polyacrylamide SDS-PAGE gel and analyzed by autoradiography.

**Western blotting of phosphorylated Smad2.** HaCat and GPI anchor mutated cells were grown in 60 mm dishes (Falcon) to 70-80% confluency and serum starved overnight. Cells were washed with PBS and were treated with various doses of TGF- $\beta$ 1 or TGF- $\beta$ 2 for the indicated times at 37°C. The cells were solubilized with 500  $\mu$ l of lysis buffer for at least 30 minutes at 4°C with mild agitation. The cell lysates were collected and centrifuged for 10 minutes at 12 000 x g. The protein concentration of the each sample was normalized to 50  $\mu$ g using the BioRad protein assay kit as recommended by the manufacturers. 1/5 volume of 5X electrophoresis reducing sample buffer was added to the sample and boiled for 5 minutes. The samples were separated on a 7.5% SDS-polyacrylamide gel and transferred to a nitrocellulose membrane. The membrane was blocked for at least three hours to

overnight in blocking buffer [TBS-T buffer: 30 mM Tris (pH 7.5), 150 mM NaCl , 0.05% w/v Tween 20 , containing 5% w/v of skim milk powder]. The blot was then incubated for 60 minutes at room temperature with the PS2 antibody (1:2000 in TBS-T buffer) which recognizes the phosphorylated C-terminus of Smad2 (a gift from Dr. S. Soucheinytskyi, Uppsala, Sweden). The blot was then washed in TBS-T buffer and incubated for 45 minutes at room temperature with a secondary anti-rabbit antibody conjugated to horseradish peroxidase (HRP) (1:5000) (Pierce, Rockford, IL.). The membrane was then subjected to chemiluminescence analysis (ECL) as detailed by the manufacturers (Amersham-Pharmacia-Biotech). In order to determine equal protein loading, membranes were immunoblotted with a goat polyclonal anti-Smad2 antibody (Santa Cruz) that is raised against a peptide near the N-terminus, or a rabbit polyclonal anti-STAT3 antibody (Santa Cruz) which recognizes a peptide mapping at the C-terminus.

## RESULTS

### 15 Isolation and cloning of a cell line mutated in GPI anchor biosynthesis:

HaCat cells were treated with 300 µg/ml of ethylmethanesulfonate (EMS) for 24 hours as described in "Materials and Methods.". EMS is an ethylating agent that has been used to cause mutagenesis of genes, including those involved in the biosynthesis of the GPI anchor (Sega, 1984; Stevens, 1999). Because the identity of the r150 is not known, another GPI-anchored protein that is strongly expressed in human keratinocytes was used in the cloning of GPI anchor deficient cells. CD59 is a 21 kDa GPI-anchored protein that is expressed in keratinocytes and is an inhibitor of the membrane attack complex (Venneker et al, 1994; Pasch et al, 1998). After EMS treatment, cells were stained with an anti-CD59 antibody conjugated to FITC and negatively sorted. Cells selected into the CD59 negative population were cloned using the FACS Vantage automated cell deposition unit. Ten clones survived and were reanalyzed for CD59 expression by flow cytometry. Two clones displayed decreased expression of CD59 on their cell surface and were presumed to be mutated in GPI anchor biosynthesis (termed GPI M and GPI M1). Figure 6A demonstrates the flow cytometric analysis of parental HaCat cells as compared to GPI M cells. Though the expression of CD59 was not



completely abolished in the GPI M cells, the immunofluorescence intensity of CD59 was analyzed to be approximately half that of HaCat cells (56.3%  $\pm$  7.4,  $p < 0.0006$ ). Similar results were obtained with the GPI M1 cells.

5 In order to ascertain that the GPI anchor mutated cells are deficient in r150, affinity labeling with  $^{125}\text{I}$ -TGF- $\beta$ 1 was performed. As demonstrated in Figure 6B, GPI M cells exhibit a significant loss of r150 on their cell surface as compared to HaCat cells. The expression of the types I, II and III receptors and their binding to  $^{125}\text{I}$ -TGF- $\beta$ 1 are unaffected in the GPI M cells and appear identical to that of parental HaCat cells.

10 **Cell growth and morphology are unchanged in GPI anchor mutated cells:**

A deficiency or loss of the cell surface expression of GPI-anchored proteins as a result of EMS mutagenesis is reported not to affect cell morphology or cell cycle progression (Stevens, 1999). In order to confirm that cell growth was not altered in the GPI anchor deficient keratinocyte cell line, the doubling time of these cells was determined. As seen in Figure 7A, GPI M cells display a similar doubling time (approximately 16hrs.) as the parental HaCat cells. In addition, the cellular morphology of GPI M cells (Figure 7C) is unchanged to that of the HaCat (Figure 7B).

**r150 negatively modulates TGF- $\beta$  stimulated transcription:**

20 TGF- $\beta$ 1 induced transactivation of the p3TP-Lux reporter construct in GPI M and HaCat cells was then evaluated. As demonstrated in Figure 8A, upon treatment with 10 pM TGF- $\beta$ 1 for 4 hours, the GPI M cells display an approximately 30% higher fold increase of luciferase activity, as compared to HaCat cells ( $p < 0.05$ ). In response to 100 pM TGF- $\beta$ 1, the difference between the GPI M and HaCat cells rises to 50% ( $p < 0.003$ ). Interestingly, the fold induction in the HaCat cells with 10 pM and 100 pM TGF- $\beta$ 1 treatment remain similar to each other (3.3 and 3.9 respectively), while the GPI M cells display a dose response, demonstrating fold increases of 4.9 and 8.4 respectively.

- 40 -

Upon 16 hours of TGF- $\beta$ 1 treatment, the GPI M cells display a markedly enhanced response to 10 pM and 100 pM TGF- $\beta$ 1 compared to the parental HaCat (Figure 3B). Also, a dose response to the 10 pM and 100 pM TGF- $\beta$ 1 are evident for both cell types. A clone of HaCat cells that was subjected to the same cloning procedure as the GPI M cells, but not mutated in GPI anchor biosynthesis (GPI NM) was included as a control. Upon 10 pM TGF- $\beta$ 1 treatment, GPI M cells display the highest fold induction of 21.4 compared to the 13.6 and 10.6 increases demonstrated by the HaCat and GPI NM cells ( $p < 0.005$  for GPI M vs HaCat). Similarly, in response to 100 pM TGF- $\beta$ 1, the fold induction exhibited by the GPI M cells is approximately twice that of HaCat and GPI NM cells (91.2 compared to 50.2 and 51.3 respectively) ( $p < 0.01$  for GPI M vs HaCat). In response to 100 pM TGF- $\beta$ 2, there is no significant difference in the stimulation of luciferase activity among the GPI M, HaCat and GPI NM cells (Figure 8C).

**GPI anchor mutated cells display enhanced Smad2 phosphorylation at low doses of TGF- $\beta$ 1:**

In order to determine if the loss of r150 had any effect on the endogenous phosphorylation of Smad2, HaCat and GPI M cells were treated with 1-50 pM of TGF- $\beta$ 1 for 20 minutes and immunoblotting was performed with an antibody raised against the phosphorylated form of Smad2. As demonstrated in Figure 9A, GPI M cells display an enhanced level of Smad2 phosphorylation as compared to the parental HaCat. In both cell types, Smad2 phosphorylation becomes detectable at 2 pM TGF- $\beta$ 1 and increases in a dose dependent fashion. However, GPI M cells exhibit an enhanced Smad2 phosphorylation in response to low (2 and 5 pM), intermediate (10 pM) and higher (50 pM) doses of TGF- $\beta$ 1 in comparison to the HaCat cells. Immunoblotting with an anti-Smad2 antibody that detects both the unphosphorylated and phosphorylated forms of Smad2 not only demonstrates equivalent protein loading, but shows that the enhanced Smad2 phosphorylation is not a result of an increased expression of total Smad2. Equal protein loading of HaCat and GPI M cell lysates is also confirmed upon immunoblotting with an anti-STAT3 antibody. Enhanced Smad2 phosphorylation in the GPI M cells is not evident

- 41 -

in response to TGF- $\beta$ 2, as HaCat and GPI M cells appear to demonstrate the same weak pattern of TGF- $\beta$ 2 induced Smad2 phosphorylation (Figure 9B).

**Elevated Smad2 phosphorylation is sustained in GPI anchor mutated cells:**

Was further examined if the GPI anchor deficient cells exhibit enhanced Smad2 phosphorylation as compared to HaCat cells upon prolonged exposure to TGF- $\beta$ 1 (Figure 10). In response to 100 pM TGF- $\beta$ 1 for the indicated times, maximal stimulation of Smad2 phosphorylation in HaCat cells is achieved at 10 minutes and maintained for 180 minutes. In GPI M cells, Smad2 is maximally phosphorylated between 45-90 minutes, followed by a moderate decrease in its phosphorylation at 180 minutes. In comparison to HaCat cells, the GPI M cells display a significantly elevated Smad2 phosphorylation which is evident at 45 minutes and is sustained until 180 minutes. The second GPI anchor mutated clone (GPI M1) also exhibits enhanced Smad2 phosphorylation in response to TGF- $\beta$ 1 (Figure 10B). In this experiment, the GPI M1 cells display an elevated level of TGF- $\beta$ 1 induced Smad2 phosphorylation as compared to HaCat cells starting at 20 minutes and is sustained until 180 minutes. On the other hand, Smad2 phosphorylation in GPI NM control cells is similar to that of HaCat cells for all the time periods studied in response to TGF- $\beta$ 1 (Figure 10C). Immunoblotting with the anti-STAT3 antibody demonstrates equal protein loading of the cell lysates. These results indicate that only the GPI anchor deficient cells (GPI M and GPI M1) exhibit enhanced TGF- $\beta$ 1 stimulated phosphorylation of endogenous Smad2 as compared to the parental HaCat cells.

**Autophosphorylation of type II kinase in HaCat and GPI anchor mutated cells.**

Since r150 can interact with the TGF- $\beta$  signaling receptors in human keratinocytes, and TGF- $\beta$  induced Smad2 phosphorylation is enhanced in GPI anchor deficient cells, we analyzed for alterations in the autophosphorylated state of the constitutively active type II receptor kinase using the *in vitro* kinase assay. As demonstrated in Figure 11, GPI M cells do not display any marked differences in type II receptor kinase phosphorylation as compared to the parental HaCat cells in either the absence or presence of 100 pM TGF- $\beta$ 1. The moderate decrease in TGF- $\beta$ RII

kinase autophosphorylation displayed by the GPI M cells, as compared to HaCat cells, did not occur in a reproducible manner.

## DISCUSSION

A keratinocyte cell line mutated in GPI anchor biosynthesis (GPI M) derived by CD59 negative based FACS from HaCat cells treated with EMS (Stevens, 1999) was created and isolated. In comparison to the parental HaCat, these cells demonstrate a significant loss in expression of the novel GPI-anchored TGF- $\beta$ 1 accessory receptor, r150, from their cell surface. There is no alteration in TGF- $\beta$  types I and II receptor expression, nor in their abilities to bind TGF- $\beta$ 1. In addition, GPI M cells do not show any differences in cellular morphology or doubling time. However, these cells display an enhanced transactivation of the TGF- $\beta$  responsive p3TP-Lux reporter gene construct as compared to HaCat cells. Furthermore, GPI M cells display an enhanced Smad2 phosphorylation in response to TGF- $\beta$ 1 treatment in a time and dose dependent manner. Taken together, the present work indicates that the novel accessory receptor, r150 acts as a negative modulator of TGF- $\beta$  action in human keratinocytes.

In the isolated GPI anchor deficient cell lines, the cell surface expression of CD59 is still detectable by flow cytometry, thus indicating that the cells do not display a complete abrogation of GPI-anchored proteins. However, a decrease in immunofluorescence intensity of approximately 50% exhibited by the GPI M cells is comparable to the loss of GPI-anchored proteins from the cell surface after PIPLC treatment (Screaton et al, 2000). In addition,  $^{125}$ I-TGF- $\beta$ 1 affinity cross-link labeling of GPI M cells demonstrate a significant loss of r150 from their cell surface as compared to HaCat cells. That GPI anchor biosynthesis is not completely abolished in these cells is likely due to the presence of GPI anchor biosynthetic genes that are resistant to EMS mutagenesis (Stevens et al, 1996). The present inventors are the first to report of the isolation of a GPI anchor deficient keratinocyte cell line.

EMS is an ethylating agent that has been used to cause mutagenesis of genes, including those involved in the biosynthesis of the GPI anchor

(Sega, 1984; Stevens, 1999). There is a mutation frequency of approximately one in a million cells upon treatment of mammalian cells with this mutagen in the dose range of 100-400 µg/ml. Hence, to "knock out" two copies of a gene, the frequency becomes 1 000 000 x 1 000 000 (Boyd and Massague, 1989; V. Stevens, personal communication). At this low frequency, the assumption is made that there is one mutation in each cell that is selected and cloned. For example, using this methodology, Stevens et al (1996) created a Chinese Hamster Ovary (CHO) cell line mutated in GPI anchor biosynthesis. This cell line has been recently used to study the role of GPI-anchored proteins in the development of Alzheimer's disease (Sambamurti et al, 1999) and in the "cross-talk" between caveolae and GPI-enriched lipid microdomains (Abrami et al, 2001). In addition, prior to the cloning of the TGF-β receptors, EMS mutagenesis was used to create the mutant Mv1Lu cell lines known as "R" and "DR" which do not express the type I and types I and II TGF-β receptors respectively (Boyd and Massague, 1989; Laiho et al, 1990). These TGF-β unresponsive R and DR mutant cell lines continue to be used to delineate components of the TGF-β signaling pathway (Massague, 1998). Therefore, the main drawback of the above model is that the expression of all GPI-anchored proteins may be affected. Thus, it is possible that the loss of other GPI proteins besides gp150 may have an impact on TGF-β signaling. However, none of the presently identified mammalian GPI-anchored proteins can bind TGF-β or are implicated in TGF-β signaling (for review, Low, 1989; Turner, 1994). The novel 150 kDa accessory receptor is the strongest candidate because it appears to be the only GPI-anchored protein that binds to TGF-β1 in keratinocytes. Furthermore, it has the potential to modulate TGF-β signaling through its interaction with the types I and II TGF-β receptors. Other GPI-anchored TGF-β binding proteins such as 180 kDa TGF-β1 binding protein and two GPI-anchored TGF-β2 binding proteins at 60 kDa and 140 kDa were identified in certain cell lines including a human osteosarcoma cell line (Cheifetz and Massague, 1991). In human fibroblasts, the present inventors identified a 180 kDa GPI-linked TGF-β1 binding protein, as well as a 65 kDa GPI-anchored TGF-β2 binding protein which do not interact with the types I and II TGF-β receptors (Dumont et al, 1995; Tam and Philip, 1998). However, early passage human keratinocytes and HaCat cells do not appear to express any of GPI-anchored TGF-β

- 44 -

binding complexes with the above mentioned relative molecular weights (Chapter 4; Tam et al, 1998). Together, it is likely that the observed differences in TGF- $\beta$ 1 induced cellular responses exhibited by the GPI M cells is due to the loss of r150.

5 The GPI anchor deficient cells display an increased transactivation of the PAI-1 promoter driven luciferase gene construct upon 4 and 16 hours of TGF- $\beta$ 1 treatment. The fold induction demonstrated by the GPI M cells is approximately twice that of HaCat cells and of an isolated keratinocyte clone that was not characterized as being mutated in GPI anchor biosynthesis (GPI NM). The conclusion that r150 is a negative modulator of TGF- $\beta$  responses is confirmed by enhanced phosphorylation of endogenous Smad2 exhibited by the GPI anchor deficient cells as compared to the parental HaCat. This phenomenon is reproducible in both GPI anchor mutants that were cloned (GPI M and GPI M1). In comparison to the HaCat cells, both GPI M and GPI M1 demonstrate a significantly elevated level of Smad2 phosphorylation which is detectable after 20-45 minutes of TGF- $\beta$ 1 treatment and sustained for as long as 180 minutes. Additionally, the GPI M cells demonstrate an enhanced Smad2 phosphorylation in response to different doses of TGF- $\beta$ 1 (2-50 pM) as compared to the parental HaCat cells. Upon treatment with TGF- $\beta$ 2, there is no detectable alteration in the pattern of Smad2 phosphorylation between HaCat and GPI M cells, indicating that the loss of r150 may not impact TGF- $\beta$ 2 responsiveness. This is also seen in the p3TP-Lux luciferase assay, whereby the HaCat and GPI M cells display similar levels of TGF- $\beta$ 2 induced luciferase activity. This is not surprising since r150 has virtually no affinity for TGF- $\beta$ 2 (Tam et al, 1998). The increased intensity and duration of the TGF- $\beta$ 1 induced Smad2 phosphorylation likely contributes to the enhanced transactivation of the PAI-1 promoter in the GPI anchor deficient cells.

r150's interaction with the types I and II receptors is reminiscent of another accessory receptor, endoglin. Endoglin is a 180 kDa homodimeric transmembrane protein that binds TGF- $\beta$ 1 and TGF- $\beta$ 3, and can interact with the types I and II receptor (see section 1.2.2 ; Cheifetz et al, 1992; Yamashita et al, 1994b). Like the r150, it possesses no kinase domain, but endoglin displays a short cytoplasmic region that is constitutively phosphorylated (Yamashita et al, 1994b).

- 45 -

Furthermore, overexpression of endoglin exerts an inhibitory effect on TGF- $\beta$  mediated growth inhibition, PAI-1 induction and angiogenesis (Letamendia et al, 1998; Li et al, 2000). However, unlike r150, endoglin cannot bind TGF- $\beta$  in the absence of the type II receptor and hence, its interaction with the signaling receptors is a ligand induced phenomenon. Furthermore, endoglin is predominantly expressed in endothelial cells while our studies indicate that r150 is not found in endothelial cell types (Wong et al, 2000; Tam and Philip, unpublished observations). Therefore, it is possible that the mechanisms by which r150 and endoglin exert their inhibitory effects are distinct from each other.

What are the potential molecular mechanisms by which r150 regulates TGF- $\beta$  signaling? Through its interaction with the TGF- $\beta$  signaling receptors, r150 may modulate the serine/threonine kinase activity of the types I or II receptors. The autophosphorylation of key residues in the type II receptor is important in the regulation of kinase function (Luo and Lodish, 1997). *In vitro* kinase assay with the anti-type II receptor antibody indicates that there is no marked difference in the type II autophosphorylation in GPI anchor deficient cells as compared to that of HaCat cells in the absence or presence of TGF- $\beta$ 1 (Figure 11). It is possible that r150 may interact with the type I receptor to hamper its phosphorylation by the type II receptor, and hence, activation of the type I kinase. This would be similar to the action of FKBP12, an immunophilin which interacts with the type I kinase and exerts an inhibitory effect on TGF- $\beta$  mediated cellular responses (Wang et al, 1996). However, our attempts to assess the phosphorylation of the type I kinase by the *in vitro* kinase assay with the anti-type I receptor antibody using previously described methods were unsuccessful (Wrana et al, 1994; Wieser et al, 1995). Alternatively, r150's interaction with the type II receptor may exert a negative regulatory effect on TGF- $\beta$  signaling as is seen with TRIP-1, a WD domain protein that specifically associates with the type II receptor and suppresses TGF- $\beta$  induced transactivation of the PAI-1 promoter (Choy and Derynck, 1998). However, it is presently unclear if r150 interacts with the signaling receptors in the absence of TGF- $\beta$ , or if the interaction is a strictly a ligand induced phenomenon. In addition, it is unknown if r150 preferentially interacts with the type I or type II receptor.

Our previous work indicates that soluble r150 retains its ability to bind TGF- $\beta$ 1 in absence of the types I and II receptors and of an intact membrane (Tam et al, 2001). Hence, it is likely that r150 binds to TGF- $\beta$ 1 independently from the TGF- $\beta$  signaling receptors when attached to the cell surface. This is in contrast to the type I receptor kinase and endoglin which only recognize TGF- $\beta$  bound to the type II receptor (Cheifetz et al, 1992; Wrana et al, 1994; Letamendia, 1998). Therefore, the membrane-anchored form of r150 may regulate TGF- $\beta$ 1 binding to its receptors. This possibility can be interpreted from the results which show that GPI anchor deficient cells display an enhanced Smad2 phosphorylation even at low and intermediate doses of TGF- $\beta$ 1 (2-10pM) as compared to the parental HaCat. Furthermore, in the p3TP-Lux luciferase assay, a dose response is demonstrated by the mutant cells upon 4 hours of 10 pM (4.9 fold induction) and 100 pM TGF- $\beta$ 1 (8.4 fold induction) treatment. In contrast, the fold increases in HaCat cells are similar at both doses (3.3 and 3.9 respectively). Hence, the access of TGF- $\beta$ 1 to its receptors appears to be ameliorated in the absence of r150. Therefore, unlike the type III receptor whose role is to facilitate binding of TGF- $\beta$  to its signaling receptors, r150 may not play the role as an "enhancer" of TGF- $\beta$  binding (Lopez-Casillas et al, 1993). As a result, the membrane-anchored r150 may act to sequester TGF- $\beta$ 1, but not TGF- $\beta$ 2 or TGF- $\beta$ 3, away from the signaling receptors on the cell surface. Thus, in the absence of r150, lower amounts of TGF- $\beta$ 1 can induce the heteromerization of the types I and II receptors which can surpass the signaling threshold required to activate the type I kinase, leading to Smad2 phosphorylation.

r150 may potentially have a role in the cellular "compartmentalization" of the types I and II receptors and of the R-Smads. There is an emerging theme in signal transduction biology whereby signaling molecules can be compartmentalized into membrane entities known as caveolae and "lipid rafts." These are organized lipid microdomains which serve as centres in which signaling molecules of various pathways can effectively interact (Sargiacomo et al, 1993; Horejsi et al, 1999). GPI-anchored proteins have been detected in caveolae or lipid rafts in association with other signaling molecules such as Src-like kinases, G-proteins, PKC, and PDGF receptor (Sargiacomo et al, 1993; Lisanti et al, 1994; Oka et al, 1997; Liu et al, 1997). The association of r150, a GPI-anchored protein, with



th type I and II receptors suggests that the TGF- $\beta$  receptors may also be organized in these plasmalemmal entities. This is a possibility since caveolin-1, the main scaffolding protein in caveolae, was recently reported to co-localize with the type I and II receptors and Smad2 in caveolae-enriched membrane fractions. The type I receptor was demonstrated to directly associate with caveolin-1 which resulted in the downregulation of TGF- $\beta$  mediated transcriptional responses and suppression of ligand induced Smad2 phosphorylation (Razani et al, 2001). Furthermore, the FYVE domain in Smad anchor for receptor activation (SARA), a protein that plays a prominent role in controlling the subcellular localization of Smad2, was required to maintain SARA localized in punctate "spots" near the cell surface as characterized by immunofluorescence microscopy (Tsukazaki et al, 1998). SARA was shown to recruit Smad2 into these domains where TGF- $\beta$  receptors also co-localized. The identity of these "spots" were not confirmed, but are speculated to be lipid rafts. Presently, it is not known whether SARA interacts with r150. That SARA and r150 may synergistically act to localize r150 into lipid microdomains is an interesting possibility. Within caveolae or lipid rafts, the TGF- $\beta$  signaling machinery may also potentially interact with signaling molecules of other pathways that reside in these structures. This would provide a structural explanation as to how TGF- $\beta$  can elicit the participation of multiple signaling pathways.

GPI-anchored proteins are implicated in the maintenance of skin homeostasis. Targeted deletion of the GPI anchor biosynthetic gene, PIG-A, in the epidermis of transgenic mice results in smaller pups exhibiting skin with a wrinkled appearance and thickened stratum corneum as compared to wild type mice (Tarutani et al, 1997). This defect in skin development is the apparent cause of death of these mice within 1-3 days after birth. Interestingly, transgenic mice engineered to overexpress TGF- $\beta$ 1 in the epidermis also demonstrate a compact stratum corneum with an increase in the stratified cell layers as compared to wild type mice (Sellheyer et al, 1993). Due to the overexpression of TGF- $\beta$ 1, there is a significant reduction in the number of proliferating epidermal cells as determined by pulse labelling with 5-bromodeoxyuridine. Death of these transgenic mice is also attributed to abnormal skin development. Taken together, the targeted overexpression of TGF- $\beta$ 1 in the epidermis appears to mimic that of the PIG-A deletion. It is possible to

envision that the ablation of r150's expression in the epidermis as a consequence of the PI3-A deletion would result in the loss of its inhibitory effects in TGF- $\beta$  action, thus contributing to the subsequent hyperactivity of TGF- $\beta$ 1 signaling in the skin. Hence, r150 may play an essential role in skin development as a key regulator of TGF- $\beta$ 's function in epidermal differentiation and homeostasis.

In conclusion, studies of TGF- $\beta$  responses in the GPI anchor deficient keratinocytes have provided critical insight into r150's function in TGF- $\beta$  signaling. The present results indicate that r150 negatively modulates TGF- $\beta$  action in human keratinocytes. In r150 deficient keratinocytes, TGF- $\beta$  induced Smad2 phosphorylation and PAI-1 expression are enhanced. It is conceivable that r150 may directly modulate type I and II kinase activity through its interaction with the signaling receptors. In addition, due to r150's ability to bind TGF- $\beta$ 1 on its own, the membrane bound and soluble r150 may regulate ligand availability by acting as a scavenger receptor. Delineation of r150's structure is necessary to elucidate the precise molecular mechanisms by which this novel accessory receptor regulates TGF- $\beta$  signaling.

### EXAMPLE 3

#### SEQUENCE OF r150 AND NUCLEIC ACIDS ENCODING THE SAME

**Cloning of r150 and expression of r150 gene confirm that r150 is an inhibitor of TGF- responses:** To determine its structural identity, r150 was purified on a TGF- $\beta$ 1 affinity column and analyzed by tandem mass spectrometry (Harvard Microchemistry Facility, Harvard Univ) which allowed us to obtain a 19 amino acid microsequence. This matched to the 5' end of an expressed sequence tagged (EST) cDNA clone of unknown function from human placenta. Subsequently, sequencing by PCR using seven primers, the full sequence was obtained. Alignment and conserved domain analysis revealed it to be a novel protein of 1428 amino acids with an N-terminal signal sequence and a C-terminal GPI anchor attachment signal sequence (Figure 12 and SEQ ID No: 2). The most likely GPI anchor cleavage site ( $\omega$ ) is at amino acid residue 1404. The predicted molecular mass and structural features indicate that this gene product represents r150. This novel TGF- accessory receptor has a thiolester signature motif and belongs to the complement C3/ 2-

macroglobulin superfamily as recently found in Lin *et al.* (2002). These authors, using a very different strategy (affinity binding of blood cells types with monoclonal antibodies) have identified a protein which has 1445 amino acids, which is very similar to r150. No definite function has been assigned to this protein called CD109, considered by the present inventors to be a variant of r150. Expression studies in HaCaT and 293 cells confirm that the cDNA encodes a 150 kDa GPI- anchored protein.

Transfection of HaCaT and 293 cells with the r150 gene (SEQ ID NO: 1) and detection of the expressed protein by an antibody specific for GPI anchor (anti-CRD antibody) demonstrate that the expressed protein migrates at 150 kDa and that it contains a GPI anchor as expected of r150 (Figure 13). Furthermore, overexpression of r150 in HaCaT and 293 cells results in strong negative modulation of TGF- $\beta$  induced Smad 2 phosphorylation and gene promoter activity, providing strong evidence to confirm that r150 is an inhibitor of TGF- $\beta$  signaling and responses. Cells transfected with r150 displayed markedly decreased Smad 2 phosphorylation upon stimulation with TGF- $\beta$  when compared with untransfected cells or cells transfected with the empty vector (Figure 14). Similarly, transfection of cells with the 3TPLUX promoter-reporter construct (encoding plasminogen activator inhibitor promoter linked to the luciferase reporter gene) resulted in a marked decrease in both the basal and TGF- $\beta$  induced promoter activity (Figure 15). Taken together, our results demonstrate that r150 is an inhibitor of TGF- $\beta$  signaling and responses *in vitro* and implicate r150, in its membrane anchored and/or soluble form, as a key regulator of TGF- $\beta$  action *in vivo* (Figure 16).

Sequence comparisons with the ones disclosed by Lin *et al.* (2002) and Schuh *et al.* (2002) provide indications of variants (see Figure 17). Some amino acid residues may change, but would presumably not change the property of binding TGF- $\beta$ 1. At least two types of r150-like would exist: one with Tyr, one with Ser at position 703. The TGF- $\beta$ 1 binding region can be predicted by sequence comparison with the sequences disclosed for  $\alpha_2$  macroglobulin (Webb *et al.* 1998). The TGF- $\beta$ 1 binding domain is ascribed between amino acids 591 and 774 of  $\alpha_2$ -M protein sequence. The minimal binding sequence appears to be the 16-mer

WDLVVUNSAGVAEVGU (Webb et al. 2000). r150 corresponding sequence shows little homology with this sequence. However, the surrounding sequences are more homologous to r150. Besides that, r150 appears to be much more selective than  $\alpha_2M$  for TGF- $\beta$ 1. It is therefore possible that the above 16-mer is sort of a "consensus" sequence which allows versatility in the binding of a plurality of cytokins. The corresponding r150 sequence would be much more specific to TGF- $\beta$ 1. This sequence is SEQ ID No: 12 for r150 and its coding nucleic acid is defined in SEQ ID NO: 11. Any protein comprising this particular minimal 16-mer is within the scope of this invention.

#### 10 Example 4:

##### Mapping the TGF- $\beta$ binding domain of r150

To confirm that r150 binding domain corresponds to the one found for  $\alpha_2M$ , the following procedure is performed.

The ligand binding domain of r150 is mapped by producing deletion mutants of r150 and analyzing ligand binding activity. The requirement for the GPI anchor for r150 function is examined using chimeric constructs in which the C-terminal GPI anchor sequence of r150 is replaced with the transmembrane domain (TM) or the GPI anchor sequence of an irrelevant protein, and determining alterations in TGF- $\beta$  responses. The deletion mutagenesis and creation of chimeric constructs is done as described previously in collaboration with Dr. Uri Saragovi, McGill. (Taheri et al, 2000; Zaccaro et al, 2001)

Deletion mutants are generated using progressive digestion of r150 cDNA using BAL-31 exonuclease starting ~20 amino acids downstream from the signal peptide. Digests are repaired and the fragments are sub-cloned to reattach the signal peptide. The deletion mutants, the full length r150 or the empty vector are expressed in COS-7 cells and in GPI mutant keratinocytes as previously described (Pepin et al, 1994). Cells will then be affinity labeled with  $^{125}I$ -TGF- $\beta$  1 and analyzed by SDS-PAGE to determine ligand binding activity as described previously (Tam and Philip, 1998). The generation of chimeric constructs (r150 in which its GPI

- 51 -

anchor sequence is replaced with the TM of insulin receptor or with the GPI anchor sequence of NCAM), was performed using overlapping PCR as previously described (Screaton et al, 2000; Zaccaro et al, 2001). Wild type r150 and the chimeric constructs are expressed in COS-7 cells and in GPI mutant keratinocytes, and TGF- $\beta$  responses are determined.

5

Although the present invention has been described hereinabove by way of preferred embodiments thereof, it can be modified, without departing from the spirit and nature of the subject invention as defined in the appended claims.

## REFERENCES

- Blobe, GC., Schiemann, WP., Lodish, H. (2000). Role of Transforming growth factor- $\beta$  in human disease. *NEJM*. 342: 1358.
- 5 Bordier C. 1981. Phase separation of integral membrane proteins in Triton X-114 solution. *J Biol Chem* 256: 1604-1607.
- Boyd FT., and Massague J. (1989). Transforming growth factor- $\beta$  inhibition of epithelial cell proliferation linked to the expression of a 53 kDa Membrane receptor. *J. Biol. Chem.* 264: 2272-2278.
- 10 Broomfield SJ, Hooper NM. 1993. Characterization of an antibody to the cross-reacting determinant of the GPI-anchor of human membran dipeptidase. *Biochim Biophys Acta* 1145: 212-218.
- Brown D. 1993. The tyrosine kinase connection: how GPI-anchored proteins activate T cells. *Curr Opinion Immunol* 5: 349-354.
- 15 Brown DA, London E. 1998. Functions of lipid rafts in biological membranes. *Ann Rev Cell Develop Biol* 14: 111-136.
- Chajek-Shaul T, Halimi O, Ben-Naim M, Stein O, Stein Y. 1989. Phosphatidylinositol-specific phospholipase releases lipoprotein lipase from the heparin releasable pool in rat heart cell cultures. *Biochim. Biophys Acta* 1014: 178-183.
- 20 Cheifetz S. and Massague J. (1991). Isoform specific transforming growth factor- $\beta$  binding proteins with membrane attachments sensitive to phosphatidyl-specific phospholipase C. *J. Biol. Chem.* 266: 20767-20772.
- Cheifetz S., Bellon T., Cales C., Vera S., Ernabeau C., Massague J., and Letarte M. (1992). Endoglin is a component of the TGF- $\beta$  receptor system in human endothelial cells. *J. Biol. Chem.* 267: 19027-19030.
- 25 Chen R., Walter EI., Parker G., Lapurga JP., Millan JL., Ikehara Y., Udenfriend S., and M dorf ME. (1998). Mammalian glycosylphosphatidylinositol

anchor transfer to proteins and post-transfer deacylation. *Proc. Natl. Acad. Sci. USA* 95: 9512-9517.

Choi ME. 1999. Cloning and characterization of a naturally occurring soluble form of TGF- $\beta$  type I receptor. *Am J Physio* 276: F88-F95.

- 5 Choy L., and Derynck R. (1998). The type II TGF- $\beta$  receptor-interacting protein TRIP-1 acts as a modulator of the TGF- $\beta$  response. *J. Biol. Chem.* 273: 31455-31462.

- Clark RAF. 1996. Wound Repair: overview and general considerations. In: Clark RAF, editor. *The Molecular and Cellular Biology of Wound Repair*. New York: Plenum Press, pp.3-35.
- 10

Dumont N, O'Connor-McCourt MD, Philip A. 1995. Transforming growth factor- $\beta$  on human endometrial cells: identification of the type I, II and III receptors, and glycosyl-phosphatidyl inositol anchored TGF- $\beta$  binding proteins. *Mol Cell Endocrinol* 111: 57-66.

- 15 Germain L, Rouabhia M, Guignard R, Carrier L, Bouvard V, Auger FA. 1993. Improvement of human keratinocyte isolation and culture using thermolysin. *Burns* 19: 99-104.

- Glick AB, Kulkarni A, Tannenbaum T, Hennings H, Flanders KC, O'Reilly M, Sporn. MB, Karlsson S, Yuspa SH. 1993. Loss of expression of TGF- $\beta$  in skin and skin tumors is associated with hyper proliferation and high risk of malignant conversion. *Proc Natl Acad Sci USA* 90: 6076-6080.
- 20

- Glick, AB, Lee, MM, Darwiche, N, Kulkarni, AB, Karlsson, S, Yuspa, SH. 1994. Targeted deletion of TGF- $\beta$ 1 gene causes rapid progression to squamous cell carcinoma. *Genes and Dev* 8: 2429-2440.

- 25 Gougos A, Letarte M. 1990. Primary structure of endoglin, an RGD-containing glycoprotein of human endothelial cells. *J Biol Chem* 265: 8361-8364.

Grainger DJ, Metcalfe JC. 1995. A pivotal role for TGF-beta in atherogenesis. *Biol Rev* 70: 571-596.

5 Hebda PA. 1988. Stimulatory effects of TGF-beta and EGF on epidermal cell outgrowth and porcine skin explant cultures. *J. Invest, Dermatol.* 91: 440-445.

Heldin CH, Miyazono K, ten Dijke P. 1997. TGF-beta signaling from cell membrane to nucleus through SMAD proteins. *Nature* 390: 465-471.

10 Hooper NM. 1992. Identification of glycosyl-phosphatidylinositol anchor on membrane proteins. In Hooper NM and Turner AJ, editors. *Lipid Modification of Proteins: A Practical Approach*. New York: IRL Press. pp. 89-115.

Hooper NM. 1999. Detergent-insoluble glycosphingolipid/cholesterol-rich membrane domains, lipid rafts and caveolae. *Mol Memb Biol* 16: 145-156.

15 Kingsley DM (1994). The TGF-beta superfamily: new members, new receptors, and new genetic tests of function in different organisms. *Genes & Devel.* 8: 133-146.

Laiho M., Weis FMB., and Massague J. (1990). Concomitant loss of TGF-beta receptor types I and II in TGF-beta resistant cell mutants implicates both receptor types in signal transduction. *J. Biol. Chem.* 265: 18518-18524.

20 Lefer AM, Ma X-L, Weyrich AS, Scalia, R. (1993). Mechanism of the cardioprotective effect of TGF-beta 1 in feline myocardial ischemia and reperfusion. *Proc Natl Acad Sci USA* 90: 1018-1022.

25 Letamendia A, Lastres P, Botella LM, Raab U, Langa C, Velasco B, Attisano L. And Bernabeu C. (1998). Role of Endoglin in cellular responses to TGF-beta. *J. Biol. Chem.* 273: 33011-33019.

Letterio JJ, Roberts AB. 1998. Regulation of immune responses by TGF-beta. *Ann Rev Immuno* 16: 137-161.



Li C., Hampson IN., Hampson L., Kumar P., Bernab u C., and Kumar S. (2000). CD10<sup>5</sup> antagonizes the inhibitory signaling of TGF- $\beta$ 1 on human vascular endothelial cells. FASEB J. 14: 55- 64.

5 Li Chengang, Hampson IN, Hampson L, Kumar P, Bernabeu C, Kumar S. 2000. CD105 antagonizes the inhibitory signaling of transforming growth factor- $\beta$ 1 on human vascular endothelial cells. FASEB J. 14: 55-64.

Lisanti MP, Sargiacomo M, Graeve L, Saltiel AIR, Rodriguez-Boulau E. 1988. Polarized apical distribution of glycosyl phosphatidylinositol-anchored proteins in renal epithelial cell lines. Proc Natl Acad Sci USA 85: 9557-10 9561.

Lisanti MP., Scherer PE., Vidugiriene J., Tang ZL., Hermanowski-Vosatka A., Tu YH., Cook RF., and Sargiacomo M. (1994). Characterization of caveolin-rich membrane domains isolated from an endothelial rich source- implications for human disease. J. Cell. Biol. 126:111-126.

15 Liu J., Oh P., Horner T., Rogers RA., and Schnitzler JE. (1997). Organized endothelial cell surface signal transduction distinct from GPI-anchored protein microdomains. J. Biol. Chem 272: 7211-7222

Lopez-Casillas F, Payne HM, Andres JL, Massague J 1994. Betaglycan can act as a dual modulator of TGF- $\beta$  access to signaling receptors: 20 mapping of ligand binding and GAG attachment sites. J Cell Biol 124: 557-568.

Lopez-Casillas F, Wrana JL, Massague J 1993. Betaglycan presents ligand to the TGF- $\beta$  signaling receptor. Cell 73: 1435-1444.

Lopez-Casillas, F., Wrana, J.L. and Massague, J. (1993). Betaglycan presents ligand to the TGF- $\beta$  signaling receptor. Cell 73: 1435-1444.

25 Low MG. (1989). Glycosylphosphatidylinositol: a versatile anchor for cell surface proteins. FASEB J. 3: 1600-1608.

Luo K., and Lodish HF. (1997). Positive and negative regulation of type II TGF- $\beta$  receptor signal transduction by autophosphorylation on multiple serine residues EMBO J. 8: 1970-1997.

Massague J. (1998). TGF- $\beta$  signal transduction. Annu. Rev. Biochem. 67: 753-791.

McNeil H, Williams C, Guan J, Dargunow M, Lawlor P, Sirimanne E, Nikolic K, Gluckman P. 1994. Neuronal rescue with transforming growth factor-beta 1 after hypoxic-ischaemic brain injury. Neuroreport 5: 901-904.

Mehta, JL, Yang BC, Strates, BS, Mehta P. 1999. Role of TGF-beta 1 in platelet-mediated cardioprotection during ischemia-reperfusion in isolated rat hearts. Growth Factors 16: 179-190.

Metz CN, Brunner G, Choi-Muir NH, Nguyen H, Gabrilove J, Caras IW, Altszuler N, Rifkin DB, Wilson EL, Davitz MA. 1994. Release of GPI-anchored membrane proteins by a cell-associated GPI-specific phospholipase D. EMBO 13: 1741-1751.

Moulin V, Auger FA, O'Connor-McCourt M, Germain L. 1997. Fetal and postnatal sera differentially modulate human dermal fibroblast phenotypic and functional features in vitro. J Cellul Physiol 171:1-10.

Movahedi S, Hooper NM. 1997. Insulin stimulates the release of the GPI-anchored membrane dipeptidase from 3T3-L1 adipocytes through the action of a phospholipase C. Biochem J 326: 531-537.

Nosjean O., Briolay, A. and Roux, B. (1997). Mammalian GPI proteins: sorting, membrane residence and functions. Biochim. Biophys. Acta. 1331: 153-186.

Nunes I, Kojima S, Rifkin DB. 1996. Effects of endogenously activated TGF- $\beta$  on growth and differentiation of retinoic acid treated HL-60 cells. Cancer Res 56: 495-499.

O'Kane S, Ferguson MWJ. 1997. Transforming growth factor- $\beta$ s and wound healing. *Int J Biochem Cell* 29: 63-78.

Oka N., Yamamoto M., Schwencke C., Kawabe J., Ebina T., Ohno S. and Couet J., Lisanti, MP., and Ishikawa, Y. (1997). Caveolin interaction with protein kinase C. *J. Biol. Chem.* 272: 33416-33421.

Onichtchouk D, Chen Y-G, Dosch R, Gawantka V, Dellus H, Massague J, Niehrs C. 1999. Silencing of TGF- $\beta$  signaling by the pseudoreceptor BAMBI. *Nature* 401: 480-484.

Pasch MC., Okada N., Bos JD., and Asghar SS. (1998). Effects of UVB on the synthesis of complement proteins by keratinocytes. *J. Invest.Dematol.* 111: 683-688.

Patel BN, David S. 1997. A Novel Glycosyl phosphatidylinositol-anchored form of ceruloplasmin is expressed by mammalian astrocytes. *J Biol Chem* 272: 20185-20190.

Peltonen J, Hsiao LL, Jaakkola S, Solberg S, Aumailley M, Timpl R, Chu ML, Uitto J. 1991. Activation of collagen gene expression in keloids: co-localization of type I and type IV collagen and TGF- $\beta$ 1 mRNA. *J Invest Dermatol* 97: 240-248.

Philip A, Hannah R, O'Connor-McCourt M. 1999. Ectodomain cleavage and shedding of the type III transforming growth factor- $\beta$  receptor in lung membranes. *Eur J Biochem* 261: 618-628.

Philip A, O'Connor-McCourt MD. 1991. Interaction of TGF- $\beta$  with alpha 3-macroglobulin: role in transforming growth factor- $\beta$  clearance. *J Biol Chem* 266: 22290-22296.

Pietenpol JA, Holt JT, Stein RW, Moses HL. 1990. TGF- $\beta$ 1 suppression of c-myc gene transcription: role in inhibition of keratinocyte proliferation. *Proc Natl Acad Sci USA* 87: 3758-3762.

Razani B., Zhang XL., Bitzer M., van G r dorff G., Bottinger EP., and Lisanti MP. (2001). Caveolin-1 regulates TGF- $\beta$ /SMAD signaling through an interaction with the TGF- $\beta$  type I receptor. *J. Biol. Chem.* 276: 6727-6738.

5 Roberts AB, Sporn MB. 1990. The transforming growth factor- $\beta$ s. In Sporn MB and Roberts AB, editors. *Peptide Growth Factors and their Receptors I*. New York: Springer-Verlag. pp. 419-472.

Rodriguez- Boulan, E., Powell, SK. (1992). Polarity of epithelial and neuronal cells. *Ann. Rev. Cell Biol.* 8: 395-427.

10 Rosen CL, Lisanti MP, Salzer JL. 1992. Expression of unique sets of GPI-linked proteins by different primary neurons in vitro. *J Cell Biol* 117: 617-627.

Saltiel AR. 1996. Diverse signaling pathways in the cellular action of insulin. *Am J Physiol* 270: E375-E382.

15 Sargiacomo M., Sudol M., Tang Z., and Lisanti M. (1993). Signal transducing molecules and GPI-linked proteins form Caveolin-rich insoluble complex in MDCK cells. (1993). *J. Cell Biol.* 122: 789-807.

Satiel, AR. Diverse signalling pathways in the cellular actions of insulin. *Am.J.Physiol.* 270: E375-E385.

20 Sreter RA., DeMarte L., Draber P., and Stanners CP. (2000). The specificity for the differentiation blocking activity of carcinoembryonic antigen resides in its GPI anchor. *J. Cell. Biol.* 150: 613-625.

Sega GA. (1984). A review of the genetic effects of ethyl methanesulfonate. *Mut. Res.* 134: 113-142.

25 Sellheyer K., Bickenbach JR., Rothnagel JA., Bundman D., Longley M., Greig T., Roberts AB., and Roop DR. (1993). Inhibition of skin

development by overexpression of TGF- $\beta$ 1 in the epidermis of transgenic mice. *Proc. Natl. Acad. Sci. USA*. 90: 5237-5241.

Shukla SD. (1982). Phosphatidylinositol specific phospholipase C. *Life Sci*. 30: 1323-1355.

5 Stevens VL. (1999). Selection of mammalian cell mutants in GPI biosynthesis. In: Gelb MH (Ed.). *Protein Lipidation Protocols*. Totowa: Humana Press. pp. 13-22.

10 Stevens VL., Zhang H., and Harreman M. (1996). Isolation and characterization of a Chinese hamster ovary (CHO) mutant defective in the second step of glycosylphosphatidylinositol biosynthesis. *Biochem. J*. 313: 253-258.

Tam BYY, Germain L, Philip A. 1998. TGF- $\beta$  receptor expression on human keratinocytes: a 150 kDa GPI-anchored TGF- $\beta$  binding protein forms a heteromeric complex with the types I and II receptors. *J Cell Biochem* 70: 573-586.

15 Tam BYY, Philip A. 1998. Transforming growth factor- $\beta$  receptor expression on human skin fibroblasts: dimeric complex formation of type I and type II receptors and identification of glycosyl phosphatidylinositol-anchored TGF- $\beta$  binding proteins. *J Cell Physio* 176: 553-564.

20 Tam BYY., Germain L., Hooper NM., and Philip A. Characterization of a 150 kDa accessory receptor to TGF- $\beta$  on keratinocytes: direct evidence for a GPI-anchor and ligand binding of the released form. *J. Cell. Biochem*. In press.

25 Tarutani M, Itami S, Okabe M, Ikawa M, Tezuka T, Yoshikawa K, Kinoshita T, Takeda J. 1997. Tissue-specific knockout of the mouse *Pig-a* gene reveals important roles for GPI-anchored proteins in skin development. *Proc Natl Acad Sci USA* 94: 7400-7405.

Tarutani, M., Itami, S., Okabe, M., Ikawa, M., Tezuka, T., Yoshikawa, K., Kinoshita, T., Takeda, J. (1997). Tissue specific knockout of the

- 60 -

mouse Plg-a gene rev ls important roles for GPI-anchored proteins in skin  
d v lopment. *Pro. Natl. acad. Sci. USA.* 94: 2386-2391.

Tsukazaki T., Chiang TA., Davison AF., Attisano L., and Wrana  
JL. (1998). SARA, a FYVE domain protein recruits Smad2 to the TGF- $\beta$  receptor.  
5 *Cell* 95: 779-791.

Turner AJ (1994). PIG-tailed membrane proteins. *Essays in  
Biochem.* 28: 113-127.

Venneker GT., Das PK., Das PK., Meinardi MMHM., Marle J.,  
Veen HA., Bos JD., and Asghar SS. (1994). GPI-anchored membrane proteins are  
10 constitutively down-regulated in psoriatic skin. *J. Pathol.* 172: 189-197.

Venneker, GT., Das, PK., Meinardi, MM., von Marle, J., van  
Veen, HA., Bos, JD., Asghar, SS. (1994). Glycosylphosphatidylinositol (GPI)-  
anchored membrane proteins are constitutively down-regulated in psoriatic skin. *J.*  
*Path.* 172: 189-197

15 Wang XJ, Greenlaugh DA, Bikenbach JR, Jiang A, Bundman  
DS, Kreig T, Derynck R, Roop DR. 1997. Expression of a dominant-negative type II  
TGF- $\beta$  receptor in the epidermis of transgenic mice blocks TGF- $\beta$ -mediated growth  
inhibition. *Proc Natl Acad Sci USA.* 94: 2386-2391.

20 Wang Y., Li BY., Danielson PD., Shah PC., Rockwell S.,  
Lechleider RJ., Martin J., Manganaro T., and Donahue PK. (1996). The immunophilin  
FKBP12 functions as a common inhibitor of the TGF- $\beta$  family type I receptors. *Cell*  
86: 435-444.

25 Wieser R., Wrana JL., and Massague J. (1995). GS domain  
mutations that constitutively activate T $\beta$ R-I, the downstream signaling component in  
the TGF- $\beta$  receptor complex. *EMBO J.* 14: 2199-2208.

Wong SH., Hamel L., Chevalier S., and Phillip A. (2000).  
Endoglin expression on human microvascular endothelial cells: association with

betaglycan and formation of higher order complexes with TGF- $\beta$  signaling receptors. Eur. J. Biochem. 267: 5550-5560.

Wrana JL, Attisano L, Wieser R, Ventura F and Massagué J. (1994). Mechanism of activation of the TGF- $\beta$  receptor. Nature 370, 341-347.

5 Wrana, J.L., Attisano, L., Wieser, R., Ventura, F. and Massagué, J. (1994). Mechanism of activation of the TGF- $\beta$  receptor. Nature 370: 341-347.

Xie M, Sesko AM, Low M. 1993. Glycosyl phosphatidylinositol-specific phospholipase D. Am J Physiol 265: C1156-C1166

10 Yamamoto T, Noble NA, Miller DE, Border WA. 1994. Sustained expression of TGF- $\beta$ 1 underlies development of progressive kidney fibrosis. Kidney Int 45: 916-927.

15 Yamashita H., Ichijo H., Grimsby S., Moren A., ten Dijke P., and Miyazono K. (1994b). Endoglin forms a heteromeric complex with the signaling receptors for TGF- $\beta$ . J. Biol. Chem. 269: 1995-2001.

Zambruno G, Marchisio PC, Marconi A, Vaschieri C, Mechiorri A, Giannetti A, DeLuca M. 1995. TGF- $\beta$ 1 modulates  $\beta$ 1 and  $\beta$ 5 integrin receptors and induces the de novo expression of the  $\alpha$ v $\beta$ 6 heterodimer in normal human keratinocytes for wound healing. J Cell Biol 129: 853-865.

20 Zamze SE, Ferguson MAJ, Collins R, Dwek RA, Rademacher TW. 1988. Characterization of the cross-reacting determinant (CRD) of the glycosylphosphatidyl membrane anchor of Trypanosoma brucei variant surface glycoprotein. Eur J Biochem 176: 527-534.

25 Pepin M-C. et al (1994). Mapping of the ligand binding domain of the transforming growth factor- $\beta$  receptor type III by deletion mutagenesis. Proc. Natl. Acad. Sci. 91: 6997-7001.

- 62 -

Zaccaro, M.C. et al (2001). P75 coreceptors regulate ligand dependent and ligand independent TrK receptor activation, in part by altering Trk docking subdomains. J. Biol. Chem. 276: 31023-31029.

5 Taheri, M. et al (2000). Self recognition in the Ig superfamily. Identification of precise subdomains in carcinoembryonic antigen required for intracellular adhesion. J. Biol Chem. 275: 26935-26943.

Screaton, R.A. et al (2000). Specificity for the differentiation blocking activity of carcinoembryonic antigen resides in its glycosylphosphatidyl-inositol anchor. J. cell. Biol. 150: 613-625.



**WHAT IS CLAIMED IS:**

1. The use of a protein comprising any one of the following sequences in the making of a medication for inhibiting TGF-  $\beta$ 1 activity in a biological tissue: SEQ ID Nos: 2 , 4, 6, 8, 10 and 12.
- 5 2. The use of an antagonist to a protein comprising any one of the following sequences in the making of a medication for increasing TGF- $\beta$ 1 activity in a biological tissue: SEQ ID Nos: 2 ,4, 6, 8, 10 and 12.
3. The use of a nucleic acid encoding a protein comprising any one of the following sequences in the making of medication for inhibiting TGF- $\beta$ 1  
10 activity in a biological tissue: SEQ ID Nos: 1, 3, 5, 7, 9 and 11.
4. The use of a molecule which silences the expression of a nucleic acid encoding a protein comprising any one of the following sequences in the making of medication for increasing TGF- $\beta$ 1 activity in a biological tissue: SEQ ID Nos: 1, 3, 5, 7, 9 and 11.
- 15 5. The use of claim 4 wherein said molecule is an antisense nucleic acid.
6. An isolated nucleic acid encoding a protein comprising any one of the following sequences : SEQ ID Nos: 2,4, 6, 8 and 10.
7. The nucleic acid of claim 6, which comprises any one of the  
20 following nucleotide sequences: SEQ ID No: 1 , 3, 5, 7, 9 and 11.
8. The nucleic acid of claim 6, which comprises the nucleotide sequence defined in SEQ ID No: 1, 3, 5 or 7.
9. The nucleic acid of claim 6, 7 or 8, which is a recombinant vector.
- 25 10. The nucleic acid of claim 9, which is an expression vector.
11. A recombinant host cell which comprises the nucleic acid of any

one of claims 6 to 10.

12. A method of producing a protein comprising any one of the following sequences : SEQ ID NOs: 2, 4, 6, 8, 10 and 12, which comprises the steps of:

- 5
- growing recombinant host cell as defined in claim 11 in a culture medium supporting cell growth and expression of said nucleic acid:
  - recovering the protein from the culture medium or from the cell.

10 13. A nucleic acid which is an antisense nucleic acid of any one of the nucleotide sequences defined in SEQ ID No: 1 or 5.

14. The nucleic acid of claim 13, which is a recombinant vector.

15. The nucleic acid of claim 14, which is an expression vector.

16. A recombinant host cell which comprises the nucleic acid of any  
15 one of claims 13, 14 or 15.

17. A TGF- $\beta$ 1 binding reagent, which comprises a protein comprising any one of the following sequences: SEQ ID Nos: 2, 4, 6, 8, 10 and 12.

18. A composition of matter which comprises the reagent of claim  
17, and a carrier.

20 19. The composition of claim 18, wherein said carrier is a pharmaceutical carrier.

20. The reagent as defined in claim 17, 18 or 19, wherein said protein is hydrosoluble and has the amino acid sequence of SEQ ID No: 4 or 8.

1/24




FIGURE 1

2/24




FIGURE 2

3/24




FIGURE 3




FIGURE 4

5/24




FIGURE 5

6/24




FIGURE 6 A



7/24




FIGURE 6B

8/24




FIGURE 7 A

9/24




FIGURE 7 B




FIGURE 7 C

10/24




FIGURE 8 A




FIGURE 8 B

11/24




FIGURE 9 A




FIGURE 9 B

12/24




FIGURE 10 A




FIGURE 10 B

13/24




FIGURE 10 C





FIGURE 11




Schematic diagram representing the cloned sequence of the r150 protein.

FIGURE 12



15/24



Expression of the cloned gene in HaCaT cells demonstrates that it represents r150. The expressed protein migrates at 150 kDa on SDS-PAGE and is detectable by an antibody (anti-CRD antibody) which detects GPI-anchor.

FIGURE 13

Inhibition of TGFβ signaling by r150 : Expression of r150 markedly decreases  
TGFβ-induced Smad2 phosphorylation.




FIGURE 14

17/24

Inhibition of TGF- $\beta$ -induced gene expression by r150: r150 expression markedly decreases TGF- $\beta$ 1-induced 3TPLux promoter activity.




FIGURE 15




FIGURE 16

**SUBSTITUTE SHEET (RULE 26)**

### Both membrane anchored and soluble r150 regulate TGF- $\beta$ response



20/24

1051 attgagtttttgattatactactgtcttgaagcccatctctcaacttcaagccactgtgaaggttaactgtgtgatggcaaccaactgactcttgaagaaaga  
 (351) I G P P A T T T V L L P S L A P T A T V L V T A A G A G L T L G G A  
 1156 agaaataatgtatcataacagtgcacacagagaaactatactagtactgagcggtatctaaacgtggaatcagaaatggaagctgttcagaaataaattat  
 (386) A A A V V I T V T G A A T T G T T S G S A S G A G L M G A V G L I A T  
 1261 actgtccccaaagtgaacttttaagattgaattccaactccactaaacaaagagatgaaatatataaagtgggatgccttttgagttggtggttagtggaac  
 (421) T V P G S G T P L I G P P I L G A S S G L G L L A T P L G S L S M A  
 1366 gttcatagtctgttaagtcctctagtaagacatacatccaactaaacaaagagatgaaatatataaagtgggatgccttttgagttggtggttagtggaac  
 (456) V H S L P L S P S L T T I G L L T A A G A I L V G S P P G L V V S G A  
 1471 aaagattgaaggagttaagctatatgttagtatccaggggacagttggtggcttaggaaacaaaattcaacatgttctcttaacaccagaaaattcttgg  
 (491) L A L L G L S T M V V S A G G L V A V G L G A S T M P S L T P G A S T  
 1576 actccaaaagcctgttaattgtattattgaagatgatgggaaattataadgtatgttctaaaattcctgttcagctgtttttaaataagataaag  
 (526) T P L A C V I V T T I G A A G G I I S A V L L I P V G L V P L A L I L  
 1681 ctatattggagtaagtgaaagctgaaccatctgaagagtcctcttaagatctctgtgacacagcctgactecatagttgggattgtagctgttgacaaaagt  
 (561) L T T S L V L A G P S G L V S L A I S V T G P A S I V G I V A V A L S  
 1786 gtgaatctgatgaatgcctctaatagtattacaatggaaaattgtgtccatgagttggaactttataacacagagatattatttaggcattgttcagattctttt  
 (596) V A L M A A S A A I T M G A V V H G L G L T A T G T T L G M P M A S P  
 1891 gcagctttcaggaatgtgactctgggtattgacagatgcaaacctcaagaagattatattgatgtgtttatgacaatgcagaatatgctgagaggtttatg  
 (631) A V P G G C G L T V L T A A A L T L A T I A G V T A A A G T A G A P M  
 996 gaagaaatgaaggacatatgttagatatcatgacttttcttggtaacagtcacacatgccgaagcattttccagagacttgatttggttagacaccaac  
 (666) G G A G G H I V A I H A P S L G S S P H V A L H P P G T T I L A T A  
 2101 atgggttacaggatttaccagaatttgagtaactgtacctgattctatcactcttggtggctactgttttggatctctgagacctgggtcttgacta  
 (701) M G T A I T G G P G V T V P A S I T S T V A T G P V I S G A L G L G L

FIGURE 17 (CONT.)

21/24

2206 acāactactccagtggaagctccaagcctccaaccatttttcatttttttgaatcttccctactctgttatcagaggtgaagaatttgctttggaaataactata  
 (736) T T T P V G L G A P G P P I P L A L P T S V I A G G G P A L G I T I  
 2311 ttcaattatttgaagatgccactgaggttaaggtatcatttggaadaatgacaaatttgatattctaatgacttcaaatgaataaatccacagccaccag  
 (771) P A T L L A A T G V L V I I G L S A L P A I L M T S A G I A T G H G  
 2416 cagacccttctgttccagtgaggtgggcaactgttcttttcccatcaggccacacatctgggaagaattctcatcacagtcacagctctttcaccct  
 (806) G T L L V P S G A G A T V L P P I A P T H L G G I P I T V T A L S P T  
 2521 gcttctgatgtcaccagatgatttttaaggtgaaggtgaaggtgaaggttcagatcactgaattgagatgttcttgctcttccatcaatgac  
 (841) A S A A V T G M I L V L A G G I G L S T S G S I L L A L T A A L G S  
 2626 accctgaaaacttgatgttctcctactaatcacagtgaatggcagtggaaggttcagatcactgaattgagatgttcttgctcttccatcaatgac  
 (876) T L L T L S P S P P P A T V T G S G A V G I T A I G A V L G P S I A G  
 2731 ttagctcattgattcagatgccttatggctgtggtgaacagacatgataaatttgcctccaaattttacattttggattatctgactaaagaacaaactg  
 (911) L A S L I A M P T G C G G A M I A P A P A I T I L A T L L L G L  
 2836 acagataatttgaagaaaaagctcttccatttatgggcaaggttaccagagagaacttctctatcagaggaagatggctctttcagtgcttttgggaattat  
 (946) T A A L L G L A L S P M A G G T G A G L L T G A G A G S P S A P G A T  
 2941 gaccttctggagcacttggtgtcagcttttggtttaagatgttcttgaagccgactcttacatagatatgacagaatgtgttacacagaacatacact  
 (981) A P S G S T T L S A P V L A C P L G A A P T I A I A G A V L H A T T T  
 3046 tggcttaaggacatcagaatccaacggtgaattttgggacccaggaagatgattcatagtgagcttcaagtggaataaaagtcacagtaaacattacagcc  
 (1016) T L L G H G L S A G G P T A P G A V I H S G L G G A L S P V T L T A  
 3151 tatattgtaacttctcctgggatatagaagatatacgcctaacatttgatgtgaagagctctatccatttttggagctgattcagtagaggaatttcagac  
 (1051) T I V T S L L G T A L T G P A I A V G G S I H P L G S G P S A G I S A  
 3256 aattatactcagccctataacttatgcattgtcatcagtgggaggtcctaaagcaaggaagctttgatatgtgacttgagagagagaacaagaaggtgac  
 (1086) A T T L A L I T T A L S S V G S P L A L G A L A M L T T A A G G G G G

FIGURE 17 (CONT.)

22/24

3361 atgaattctgggtgtcatcagagtcctcctcctgcagccagctcctggatattgaagttgcagctatgactgctcacacttcttaca  
 (1121) M G P T V S S G S L L S A S T G P A S L A I G V A A T A L L S H P L G  
 3466 ttccagacttctgaggaatcccaattataggttggttaagcagcaaaataagttgggtgttttgcactactcagataccactgtggcttaaggct  
 (1156) P G T S G G I P I M A T L S A G A A S L G G P A S T G A T T V A L L A  
 3571 ctgtctgaatttcagcctaatgaatacagaaggaacaatatccaaagtacccgtacgggcttagtcaccaagtcctgaaagtcttgattgacacacac  
 (1191) L S G P A A L M A T G A T A I G V T V T G P S S P S P V L P L I A T H  
 3676 aaccgttactcctcagacagcagcttctgtgtacagcaatggcagttatattccgcaaatgttttgatttgcatttgcagctcaatgttcta  
 (1226) A A L L L G T A G L A V V G P M A V A I S A A G P G P A I C G L A V V  
 3781 tataatgtgaaggcttctgggtcttctagagacgaagatctatccaaaatcaagaagccttggatttagatgttgctgtaaaagaaaataaagatgactcaat  
 (1261) T A V L A S G S A A A A S I G A G A P A L A V A V L G A L A L A  
 3886 catgtgatttgatgtgtacaagcttttcgggcccgggtgaggtggatggctcttatgaagttaacctattaagtggctttatggcgcttcagaagca  
 (1296) H V A L A V C T S P S G P G A S G M A L M G V A L L S G P M V P S G A  
 3991 atttctctgagcgagacagtgaagaaagtgaatatatcatgaaactcaacctctattagattctctaaatgaaacccagtttctgttaattctctgct  
 (1331) I S L S G T V L L V G T A H G L L A L T L A S V A G T G P C V A I P A  
 4096 gtgagaaactttaaagttcaaataccacaagatgcttcagtgatgcattagtgattactatgagccaagggagacaggggtgagaagttacaactctgaagtgaag  
 (1366) V A A P L V S A T G A A S V S I V A T T G P A A G A V A S T A S G V L  
 4201 ctgtcctcctgtgaccttgcagtgatgtccagggctccgtccttgtgaggtgagcttcaggctcccatcactcttcagtcatttttatttctgttc  
 (1401) L S S C A L C S A V G G C A P C G A G A S G S H H S S V I P I P C P  
 4306 aagcttctgtacttatggaactttggctgtga  
 (1436) L L L L T P M G L T

FIGURE 17 (CONT.)



23/24

4339                   tttatatttaaggactctgttaacactaacatttccagtagtcacatgtgattgttttgcgttagaa  
 4411                   gaatactgcttctattttgaaaaagagtttttctctctatgggttcaggagatggtgtacaacaggtcctagcatgtatagctgcatagatttcttcacc  
 4516                   tgatcttfgtgtggaagatcagaatgaatgcagttgtgtctatatttccccccacaaaatcttttagaattttttggaggtgtttgtttctccagaataa  
 4521                   aggtattactttagaa;  
 4637                   tttaggtattctctcatttttgtgaaagaaatgaacctagattctttaagcattattacacatccatgtttgtttaaagatggatttccct  
 4726                   gggaatggagaaacacgacgaggaagcttcatctgttcccttccacctccaacctagccctactgccaccccccccccaaccccccatgccagtg  
 4831                   tctcagtagatacttcttaactggaattcttcttttcagaatctaggtggtgaatttttttaagtggcacgggtcttttctgtctgaaatctcgatcacaccc  
 4936                   ccacgccattgccctccctctcttttctctgtagagaaatgtgagggcagtagacatttactgtgcttttcacacccatctcagaggttgaggagcatactgaaa  
 5041                   attgccctgggggtgctgggtgtgctgtctcttccacatcctcagccccacacccagctctatttcaggggtgagagtgcagagagcactgcaatatgtgcttc  
 5146                   atgggatttcgattcgaagatcctagaccagggagacactgttagccagggaatacaacaaaatactaggtaagtcactgcagaccgacctccctgcagtttcggga  
 5251                   aagaagctgggtttgtggagaatcagagcatcttcacatgactgctgaacctaaagatccctggcatttggccaggatcctgtggaaacctcttctagttcaggggt  
 5356                   gtgagcattagactgcaggttgtctagtacatctgatgcttgcgtgaacttttaagatccccgaatcctgagcacctcaatctttaattgacctgtattccga  
 5461                   agggtaataataatttatctggatggaaattttaagatgaatccccctttttctctctcttcttcccttctcccttctcttcttgccttctaaata  
 5566                   tactgaaatgatttagataatgtgtcaacaattaatgatcttttattcaatcgaagaaatgggttagtttttctctttagctctatggcatttcaactcaagtgagac  
 5671                   aggggaaaaagtaattgccatgggctccaaagaaattgctttatgttttagctatttaaaaataaatccatcaaaaataaagtatgcaaatgtatctttt  
 4410                   tttatatttaaggactctgttaacactaacatttccagtagtcacatgtgattgttttgcgttagaa  
 4515                   gaatactgcttctattttgaaaaagagtttttctctctatgggttcaggagatggtgtacaacaggtcctagcatgtatagctgcatagatttcttcacc  
 4620                   tgatcttfgtgtggaagatcagaatgaatgcagttgtgtctatatttccccccacaaaatcttttagaattttttggaggtgtttgtttctccagaataa  
 4636                   aggtattactttagaa;  
 4725                   tttaggtattctctcatttttgtgaaagaaatgaacctagattctttaagcattattacacatccatgtttgtttaaagatggatttccct  
 4830                   gggaatggagaaacacgacgaggaagcttcatctgttcccttccacctccaacctagccctactgccaccccccccccaaccccccatgccagtg  
 4935                   tctcagtagatacttcttaactggaattcttcttttcagaatctaggtggtgaatttttttaagtggcacgggtcttttctgtctgaaatctcgatcacaccc  
 5040                   ccacgccattgccctccctctcttttctctgtagagaaatgtgagggcagtagacatttactgtgcttttcacacccatctcagaggttgaggagcatactgaaa  
 5145                   attgccctgggggtgctgggtgtgctgtctcttccacatcctcagccccacacccagctctatttcaggggtgagagtgcagagagcactgcaatatgtgcttc  
 5250                   atgggatttcgattcgaagatcctagaccagggagacactgttagccagggaatacaacaaaatactaggtaagtcactgcagaccgacctccctgcagtttcggga  
 5355                   aagaagctgggtttgtggagaatcagagcatcttcacatgactgctgaacctaaagatccctggcatttggccaggatcctgtggaaacctcttctagttcaggggt  
 5460                   gtgagcattagactgcaggttgtctagtacatctgatgcttgcgtgaacttttaagatccccgaatcctgagcacctcaatctttaattgacctgtattccga  
 5565                   agggtaataataatttatctggatggaaattttaagatgaatccccctttttctctctcttcttcccttctcccttctcttcttgccttctaaata  
 5670                   tactgaaatgatttagataatgtgtcaacaattaatgatcttttattcaatcgaagaaatgggttagtttttctctttagctctatggcatttcaactcaagtgagac  
 5771                   aggggaaaaagtaattgccatgggctccaaagaaattgctttatgttttagctatttaaaaataaatccatcaaaaataaagtatgcaaatgtatctttt

FIGURE 17 (CONT.)

24/24

Comparison between our sequence and alpha 2 Macroglobulin  
(GI: 224053)

Identities = 63/197 (32%), Positives = 90/197 (46%), Gaps = 50/197 (25%)

```

Query: 621 YYLG-----MFMNS-----FAVFQECGLWVLTANLTKDYI 651
      + G                +++N                ++ ++ GL  T++ + K  +
Sbjct: 605 GFPGLNDQDDEDCINRHNVIYINGITYTPVSSSTNEKDMYSFLEDMGLKAFTNSKIRKPKM 664

Query: 652 DGVYDNAE-----YAERFMEENEGHIVDIHDFSLGSSPH---VRKHFPETWIWL 697
      E                Y    M          +V + +      PH  VRK+FPETWIW
Sbjct: 665 CPQLQQYEMHGPEGLRVGFYESDVMGRGHARLVHVEE-----PHTETVRKYFPETWIWD 718

Query: 698 DTNMGSRITYQEFVTVPDSTSWVATGFVISEDGLGLTTTPVELQAFQPFFFIFLNLAYS 757
      + S      E  VTVPD+IT W A F +SED GLG+++T  L+AFQPF+ L +PYS
Sbjct: 719 LVVNSAGVAEVGVTVPDITITWKGAFCLSEDAGLGISST-ASLRAFQPFVVELTMPYS 777

Query: 758 VIRGEFALETTIFNYL 775
      VIRGE F L+ T+ NYL
Sbjct: 778 VIRGEAFTLKATVLNYL 795

```

FIGURE 18

- 1 -

## SEQUENCE LISTING

<110> McGill University, through its Office of Technology Transfer  
 PHILIP, Anie  
 TAM, Betty

<120> A 150 kDa accessory receptor for TGF-beta 1

<130> DH/11168.194

<140> PCT/CA02/00560

<141> 2002-04-24

<150> US 60/285,713

<151> 2001-04-24

<150> US 60/356,163

<151> 2002-02-14

<160> 12

<170> PatentIn version 3.1

<210> 1

<211> 4369

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (2133)..(2133)

<223> N = A or C

<400> 1

tgtagcccag gcagacgccg togagatgca gggcccaccg ctctgaccg cggcccacct	60
cctctgcgtg tgcaccgccg cgctggccgt ggctcccggt cctcggtttc tggtagacagc	120
cccagggatc atcaggcccg gaggaaatgt gactattggg gtggagcttc tggaacactg	180
cccttcacag gtgactgtga aggcggagct gctcaagaca gcatcaaacc tcaactgtctc	240
tgtcctggaa gcagaaggag tctttgaaaa aggctctttt aagacactta ctcttccatc	300
actacctctg aacagtgcag atgagattta tgagctacgt gtaaccggac gtaeccagga	360
tgagatttta ttctctaata gtacccgctt atcatttgag accaagagaa tatctgtctt	420
cattcaaaca gacaaggcct tatacaagcc aaagcaagaa gtgaagtttc gcattgttac	480
actcttctca gattttaagc cttacaaaac ctctttaaac attctcatta aggaccccaa	540
atcaaatttg atccaacagt gggtgtcaca acaaagtgat cttggagtca tttccaaaac	600
ttttcagcta tcttcccatc caatacttgg tgactggctt attcaagttc aagtgaatga	660
ccagacatac tatcaatcat ttcaggtttc agaatatgta ttacaaaat ttgaagtga	720
tttgcagaca ccattatatt gttctatgaa ttctaagcat ttaaattgga ccatcacggc	780
aaagtataca tatgggaagc cagtgaagg agacgtaacg cttacatttt tacctttatc	840

cttttgggga aagaagaaaa atattacaaa aacattttaag ataaatggat ctgcaaactt 900  
 ctcttttaat gatgaagaga tgaaaaatgt aatggattct tcaaattggac tttctgaata 960  
 cctggatcta tcttcccctg gaccagtaga aattttaacc acagtgcag aatcagttac 1020  
 aggtatttca agaaatgtaa gcactaatgt gttcttcaag caacatgatt acatcattga 1080  
 gttttttgat tatactactg tcttgaagcc atctctcaac ttcacagcca ctgtgaaggt 1140  
 aactcgtgct gatggcaacc aactgactct tgaagaaaga agaaataatg tagtcataac 1200  
 agtgacacag agaaactata ctgagtactg gagcggatct aacagtggaa atcagaaaat 1260  
 ggaagctgtt cagaaaataa attatactgt ccccaaagt ggaactttta agattgaatt 1320  
 cccaatcctg gaggattcca gtgagctaca gttgaaggcc tatttccttg gtagtaaaag 1380  
 tagcatggca gttcatagtc tgtttaagtc tctagtaag acatacatcc aactaaaaac 1440  
 aagagatgaa aatataaagg tgggatcgcc ttttgagttg gtggttagtg gcaacaaacg 1500  
 attgaaggag ttaagctata tggtagtacc caggggacag ttggtggctg taggaaaaca 1560  
 aaattcaaca atgttctctt taacaccaga aaattcttgg actccaaaag cctgtgtaat 1620  
 tgtgtattat attgaagatg atggggaaat tataagtgat gttctaaaaa ttcctgttca 1680  
 gcttggtttt aaaaataaga taaagctata ttggagtaaa gtgaaagctg aaccatctga 1740  
 gaaagtctct cttaggatct ctgtgacaca gcctgactcc atagttggga ttgtagctgt 1800  
 tgacaaaagt gtgaatctga tgaatgcctc taatgatatt acaatggaaa atgtggtcca 1860  
 tgagttggaa ctttataaca caggatatta tttaggcatg ttcattgaatt cttttgcagt 1920  
 ctttcaggaa tgtggactct gggatttgac agatgcaaac ctcacgaagg attatattga 1980  
 tgggtgtttat gacaatgcag aatatgctga gaggtttatg gaggaaaatg aaggacatat 2040  
 tgtagatatt catgactttt ctttgggtag cagtccacat gtccgaaagc attttcaga 2100  
 gacttggatt tggctagaca ccaacatggg ttncaggatt taccaagaat ttgaagtaac 2160  
 tgtacctgat tctatcactt cttgggtggc tactggtttt gtgatctctg aggacctggg 2220  
 tcttgacta acaactactc cagtggagct ccaagccttc caaccatttt tcattttttt 2280  
 gaatcttccc tactctgtta tcagagggtga agaatttgct ttggaaataa ctatattcaa 2340  
 ttatttgaaa gatgccactg aggttaaggt aatcattgag aaaagtgaca aatttgatat 2400  
 tctaattgact tcaaattgaaa taaatgccac aggccaccag cagacccttc tggttcccag 2460  
 tgaggatggg gcaactgttc tttttcccat caggccaaca catctgggag aaattcctat 2520  
 cacagtcaca gctctttcac ccaactgcttc tgatgctgtc acccagatga ttttagtaaa 2580  
 ggctgaagga atagaaaaat catattcaca atccatotta ttagacttga ctgacaatag 2640  
 gctacagagt accctgaaaa ctttgagttt ctcatttcct cctaatacag tgactggcag 2700

tgaaagagtt cagatcactg caattggaga tgttcttggt ccttccatca atggcttagc 2760  
 ctcatlgatt cggatgcctt atggctgtgg tgaacagaac atgataaatt ttgctccaaa 2820  
 tatttacatt ttggattatc tgactaaaaa gaaacaactg acagataatt tgaaagaaaa 2880  
 agctctttca tttatgaggc aaggttacca gagagaactt ctctatcaga gggaagatgg 2940  
 ctctttcagt gcttttggga attatgacct ttctgggagc acttggttgt cagcttttgt 3000  
 ttttaagatgt ttccctgaag ccgatcctta catagatatt gatcagaatg tgttacacag 3060  
 aacatacact tggcttaaag gacatcagaa atccaacggg gaattttggg atccaggaag 3120  
 agtgattcat agtgagcttc aagggtggca taaaagtcca gtaacactta cagcctatat 3180  
 tgtaacttct ctctgggat atagaaagta tcagcctaac attgatgtgc aagagtctat 3240  
 ccattttttg gagtctgaat tcagtagagg aatttcagac aattatactc tagcccttat 3300  
 aacttatgca ttgtcatcag tggggagtcc taaagcgaag gaagctttga atatgctgac 3360  
 ttggagagca gaacaagaag gtggcatgca attctgggtg tcatcagagt ccaaactttc 3420  
 tgactcctgg cagccacgct ccctggatat tgaagttgca gcctatgcac tgctctcaca 3480  
 cttcttaca tttcagactt ctgagggaat cccaattatg aggtggctaa gcaggcaaag 3540  
 aaatagcttg ggtggttttg catctactca ggataccact gtggctttaa aggctctgtc 3600  
 tgaatttgca gccctaata atacagaaag gacaaatata caagtgaccg tgacggggcc 3660  
 tagctcacca agtcctcttg ctgtggtaca gccaacggca gttaatattt ccgcaaattg 3720  
 ttttggaattt gctatttgtc agctcaatgt tgtatataat gtgaaggctt ctgggtcttc 3780  
 tagaagacga agatctatcc aaaatcaaga agcctttgat ttagatgttg ctgtaaaaga 3840  
 aaataaagat gatctcaatc atgtggattt gaatgtgtgt acaagctttt cgggcccggg 3900  
 taggagtggc atggctctta tggaagttaa cctattaagt ggctttatgg tgccttcaga 3960  
 agcaatttct ctgagcgaga cagtgaagaa agtggaatat gatcatggaa aactcaacct 4020  
 ctatttagat tctgtaaatg aaacccagtt ttgtgttaat attcctgctg tgagaaactt 4080  
 taaagtttca aatacccaag atgcttcagt gtccatagtg gattactatg agccaaggag 4140  
 acaggcgggtg agaagttaca actctgaagt gaagctgtcc tcctgtgacc ttgcagtga 4200  
 tgtccagggc tgccgtcctt gtgaggatgg agcttcaggc tccatcatc actcttcagt 4260  
 catttttatt ttctgtttca agcttctgta ctttatggaa ctttggctgt gatttatttt 4320  
 taaaggactc tgtgtaacac taacatttcc agtagtcaca tgtgattgt 4369

&lt;210&gt; 2

&lt;211&gt; 1428

&lt;212&gt; PRT

&lt;213&gt; homo sapiens

- 4 -

&lt;220&gt;

&lt;221&gt; MISC FEATURE

&lt;222&gt; (703)..(703)

&lt;223&gt; Xaa = Ser or Tyr

&lt;400&gt; 2

Met Gln Gly Pro Pro Leu Leu Thr Ala Ala His Leu Leu Cys Val Cys  
 1 5 10 15

Thr Ala Ala Leu Ala Val Ala Pro Gly Pro Arg Phe Leu Val Thr Ala  
 20 25 30

Pro Gly Ile Ile Arg Pro Gly Gly Asn Val Thr Ile Gly Val Glu Leu  
 35 40 45

Leu Glu His Cys Pro Ser Gln Val Thr Val Lys Ala Glu Leu Leu Lys  
 50 55 60

Thr Ala Ser Asn Leu Thr Val Ser Val Leu Glu Ala Glu Gly Val Phe  
 65 70 75 80

Glu Lys Gly Ser Phe Lys Thr Leu Thr Leu Pro Ser Leu Pro Leu Asn  
 85 90 95

Ser Ala Asp Glu Ile Tyr Glu Leu Arg Val Thr Gly Arg Thr Gln Asp  
 100 105 110

Glu Ile Leu Phe Ser Asn Ser Thr Arg Leu Ser Phe Glu Thr Lys Arg  
 115 120 125

Ile Ser Val Phe Ile Gln Thr Asp Lys Ala Leu Tyr Lys Pro Lys Gln  
 130 135 140

Glu Val Lys Phe Arg Ile Val Thr Leu Phe Ser Asp Phe Lys Pro Tyr  
 145 150 155 160

Lys Thr Ser Leu Asn Ile Leu Ile Lys Asp Pro Lys Ser Asn Leu Ile  
 165 170 175

Gln Gln Trp Leu Ser Gln Gln Ser Asp Leu Gly Val Ile Ser Lys Thr  
 180 185 190

Phe Gln Leu Ser Ser His Pro Ile Leu Gly Asp Trp Ser Ile Gln Val  
 195 200 205

Gln Val Asn Asp Gln Thr Tyr Tyr Gln Ser Phe Gln Val Ser Glu Tyr

- 5 -

210	215	220
Val Leu Pro Lys Phe Glu Val Thr Leu Gln Thr Pro Leu Tyr Cys Ser		
225	230	235 240
Met Asn Ser Lys His Leu Asn Gly Thr Ile Thr Ala Lys Tyr Thr Tyr		
	245	250 255
Gly Lys Pro Val Lys Gly Asp Val Thr Leu Thr Phe Leu Pro Leu Ser		
	260	265 270
Phe Trp Gly Lys Lys Lys Asn Ile Thr Lys Thr Phe Lys Ile Asn Gly		
	275	280 285
Ser Ala Asn Phe Ser Phe Asn Asp Glu Glu Met Lys Asn Val Met Asp		
	290	295 300
Ser Ser Asn Gly Leu Ser Glu Tyr Leu Asp Leu Ser Ser Pro Gly Pro		
305	310	315 320
Val Glu Ile Leu Thr Thr Val Thr Glu Ser Val Thr Gly Ile Ser Arg		
	325	330 335
Asn Val Ser Thr Asn Val Phe Phe Lys Gln His Asp Tyr Ile Ile Glu		
	340	345 350
Phe Phe Asp Tyr Thr Thr Val Leu Lys Pro Ser Leu Asn Phe Thr Ala		
	355	360 365
Thr Val Lys Val Thr Arg Ala Asp Gly Asn Gln Leu Thr Leu Glu Glu		
	370	375 380
Arg Arg Asn Asn Val Val Ile Thr Val Thr Gln Arg Asn Tyr Thr Glu		
385	390	395 400
Tyr Trp Ser Gly Ser Asn Ser Gly Asn Gln Lys Met Glu Ala Val Gln		
	405	410 415
Lys Ile Asn Tyr Thr Val Pro Gln Ser Gly Thr Phe Lys Ile Glu Phe		
	420	425 430
Pro Ile Leu Glu Asp Ser Ser Glu Leu Gln Leu Lys Ala Tyr Phe Leu		
	435	440 445
Gly Ser Lys Ser Ser Met Ala Val His Ser Leu Phe Lys Ser Pro Ser		
450	455	460

- 6 -

Lys Thr Tyr Ile Gln Leu Lys Thr Arg Asp Glu Asn Ile Lys Val Gly  
 465 470 475 480

Ser Pro Phe Glu Leu Val Val Ser Gly Asn Lys Arg Leu Lys Glu Leu  
 485 490 495

Ser Tyr Met Val Val Ser Arg Gly Gln Leu Val Ala Val Gly Lys Gln  
 500 505 510

Asn Ser Thr Met Phe Ser Leu Thr Pro Glu Asn Ser Trp Thr Pro Lys  
 515 520 525

Ala Cys Val Ile Val Tyr Tyr Ile Glu Asp Asp Gly Glu Ile Ile Ser  
 530 535 540

Asp Val Leu Lys Ile Pro Val Gln Leu Val Phe Lys Asn Lys Ile Lys  
 545 550 555 560

Leu Tyr Trp Ser Lys Val Lys Ala Glu Pro Ser Glu Lys Val Ser Leu  
 565 570 575

Arg Ile Ser Val Thr Gln Pro Asp Ser Ile Val Gly Ile Val Ala Val  
 580 585 590

Asp Lys Ser Val Asn Leu Met Asn Ala Ser Asn Asp Ile Thr Met Glu  
 595 600 605

Asn Val Val His Glu Leu Glu Leu Tyr Asn Thr Gly Tyr Tyr Leu Gly  
 610 615 620

Met Phe Met Asn Ser Phe Ala Val Phe Gln Glu Cys Gly Leu Trp Val  
 625 630 635 640

Leu Thr Asp Ala Asn Leu Thr Lys Asp Tyr Ile Asp Gly Val Tyr Asp  
 645 650 655

Asn Ala Glu Tyr Ala Glu Arg Phe Met Glu Glu Asn Glu Gly His Ile  
 660 665 670

Val Asp Ile His Asp Phe Ser Leu Gly Ser Ser Pro His Val Arg Lys  
 675 680 685

His Phe Pro Glu Thr Trp Ile Trp Leu Asp Thr Asn Met Gly Xaa Arg  
 690 695 700

Ile Tyr Gln Glu Phe Glu Val Thr Val Pro Asp Ser Ile Thr Ser Trp



- 7 -

705		710		715		720
Val Ala Thr Gly Phe Val Ile Ser Glu Asp Leu Gly Leu Gly Leu Thr	725		730		735	
Thr Thr Pro Val Glu Leu Gln Ala Phe Gln Pro Phe Phe Ile Phe Leu	740		745		750	
Asn Leu Pro Tyr Ser Val Ile Arg Gly Glu Glu Phe Ala Leu Glu Ile	755		760		765	
Thr Ile Phe Asn Tyr Leu Lys Asp Ala Thr Glu Val Lys Val Ile Ile	770		775		780	
Glu Lys Ser Asp Lys Phe Asp Ile Leu Met Thr Ser Asn Glu Ile Asn	785		790		795	800
Ala Thr Gly His Gln Gln Thr Leu Leu Val Pro Ser Glu Asp Gly Ala	805		810		815	
Thr Val Leu Phe Pro Ile Arg Pro Thr His Leu Gly Glu Ile Pro Ile	820		825		830	
Thr Val Thr Ala Leu Ser Pro Thr Ala Ser Asp Ala Val Thr Gln Met	835		840		845	
Ile Leu Val Lys Ala Glu Gly Ile Glu Lys Ser Tyr Ser Gln Ser Ile	850		855		860	
Leu Leu Asp Leu Thr Asp Asn Arg Leu Gln Ser Thr Leu Lys Thr Leu	865		870		875	880
Ser Phe Ser Phe Pro Pro Asn Thr Val Thr Gly Ser Glu Arg Val Gln	885		890		895	
Ile Thr Ala Ile Gly Asp Val Leu Gly Pro Ser Ile Asn Gly Leu Ala	900		905		910	
Ser Leu Ile Arg Met Pro Tyr Gly Cys Gly Glu Gln Asn Met Ile Asn	915		920		925	
Phe Ala Pro Asn Ile Tyr Ile Leu Asp Tyr Leu Thr Lys Lys Lys Gln	930		935		940	
Leu Thr Asp Asn Leu Lys Glu Lys Ala Leu Ser Phe Met Arg Gln Gly	945		950		955	960

Tyr Gln Arg Glu Leu Leu Tyr Gln Arg Glu Asp Gly Ser Phe Ser Ala  
965 970 975

Phe Gly Asn Tyr Asp Pro Ser Gly Ser Thr Trp Leu Ser Ala Phe Val  
980 985 990

Leu Arg Cys Phe Leu Glu Ala Asp Pro Tyr Ile Asp Ile Asp Gln Asn  
995 1000 1005

Val Leu His Arg Thr Tyr Thr Trp Leu Lys Gly His Gln Lys Ser  
1010 1015 1020

Asn Gly Glu Phe Trp Asp Pro Gly Arg Val Ile His Ser Glu Leu  
1025 1030 1035

Gln Gly Gly Asn Lys Ser Pro Val Thr Leu Thr Ala Tyr Ile Val  
1040 1045 1050

Thr Ser Leu Leu Gly Tyr Arg Lys Tyr Gln Pro Asn Ile Asp Val  
1055 1060 1065

Gln Glu Ser Ile His Phe Leu Glu Ser Glu Phe Ser Arg Gly Ile  
1070 1075 1080

Ser Asp Asn Tyr Thr Leu Ala Leu Ile Thr Tyr Ala Leu Ser Ser  
1085 1090 1095

Val Gly Ser Pro Lys Ala Lys Glu Ala,Leu Asn Met Leu Thr Trp  
1100 1105 1110

Arg Ala Glu Gln Glu Gly Gly .Met Gln Phe Trp Val Ser Ser Glu  
1115 1120 1125

Ser Lys Leu Ser Asp Ser Trp Gln Pro Arg Ser Leu Asp Ile Glu  
1130 1135 1140

Val Ala Ala Tyr Ala Leu Leu Ser His Phe Leu Gln Phe Gln Thr  
1145 1150 1155

Ser Glu Gly Ile Pro Ile Met Arg Trp Leu Ser Arg Gln Arg Asn  
1160 1165 1170

Ser Leu Gly Gly Phe Ala Ser Thr Gln Asp Thr Thr Val Ala Leu  
1175 1180 1185

Lys Ala    Leu Ser Glu Phe Ala    Ala Leu Met Asn Thr    Glu Arg Thr

- 9 -

1190		1195		1200
Asn Ile Gln Val Thr Val	Thr Gly Pro Ser Ser	Pro Ser Pro Leu		
1205	1210	1215		
Ala Val Val Gln Pro Thr	Ala Val Asn Ile Ser	Ala Asn Gly Phe		
1220	1225	1230		
Gly Phe Ala Ile Cys Gln	Leu Asn Val Val Tyr	Asn Val Lys Ala		
1235	1240	1245		
Ser Gly Ser Ser Arg Arg	Arg Arg Ser Ile Gln	Asn Gln Glu Ala		
1250	1255	1260		
Phe Asp Leu Asp Val Ala	Val Lys Glu Asn Lys	Asp Asp Leu Asn		
1265	1270	1275		
His Val Asp Leu Asn Val	Cys Thr Ser Phe Ser	Gly Pro Gly Arg		
1280	1285	1290		
Ser Gly Met Ala Leu Met	Glu Val Asn Leu Leu	Ser Gly Phe Met		
1295	1300	1305		
Val Pro Ser Glu Ala Ile	Ser Leu Ser Glu Thr	Val Lys Lys Val		
1310	1315	1320		
Glu Tyr Asp His Gly Lys	Leu Asn Leu Tyr Leu	Asp Ser Val Asn		
1325	1330	1335		
Glu Thr Gln Phe Cys Val	Asn Ile Pro Ala Val	Arg Asn Phe Lys		
1340	1345	1350		
Val Ser Asn Thr Gln Asp	Ala Ser Val Ser Ile	Val Asp Tyr Tyr		
1355	1360	1365		
Glu Pro Arg Arg Gln Ala	Val Arg Ser Tyr Asn	Ser Glu Val Lys		
1370	1375	1380		
Leu Ser Ser Cys Asp Leu	Cys Ser Asp Val Gln	Gly Cys Arg Pro		
1385	1390	1395		
Cys Glu Asp Gly Ala Ser	Gly Ser His His His	Ser Ser Val Ile		
1400	1405	1410		
Phe Ile Phe Cys Phe Lys	Leu Leu Tyr Phe Met	Glu Leu Trp Leu		
1415	1420	1425		

<210> 3  
 <211> 5883  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <222> (2220)..(2220)  
 <223> N = A or C

<400> 3  
 ctaaactcga attaagaggg aaaaaaaatc agggaggagg tggcaagcca caccaccacgg 60  
 tgcccgcgaa cttccccggc agcggactgt agcccaggca gacgccgtcg agatgcaggg 120  
 cccaccgctc ctgaccgccg cccacctcct ctgcgtgtgc accgccgcgc tggccgtggc 180  
 tcccgggcct cggtttctgg tgacagcccc agggatcatc aggcccgag gaaatgtgac 240  
 tattggggtg gagcttctgg aacactgccc ttcacagggtg actgtgaagg cggagctgct 300  
 caagacagca tcaaacctca ctgtctctgt cctggaagca gaaggagtct ttgaaaaagg 360  
 ctcttttaag acacttactc ttccatcact acctctgaac agtgcagatg agatttatga 420  
 gctacgtgta accggacgta cccaggatga gattttattc tctaatagta cccgcttattc 480  
 atttgagacc aagagaatat ctgtcttcat tcaaacagac aaggccttat acaagccaaa 540  
 gcaagaagtg aagtttctga ttgttacact cttctcagat ttttaagcctt acaaaacctc 600  
 tttaaacatt ctcatthaagg accccaaatc aaatttgatc caacagtggg tgtcacaaca 660  
 aagtgatctt ggagtcattt ccaaaaacttt tcagctatct tcccatccaa tacttggtga 720  
 ctggtctatt caagttcaag tgaatgacca gacatattat caatcatttc aggtttcaga 780  
 atatgtatta ccaaaatttg aagtgaactt gcagacacca ttatattggt ctatgaattc 840  
 taagcattta aatggtacca tcacggcaaa gtatacatat gggaagccag tgaaaggaga 900  
 cgtaacgctt acatttttac ctttatcctt ttggggaaag aagaaaaata ttacaaaaac 960  
 atttaagata aatggatctg caaacttctc ttttaatgat gaagagatga aaaatgtaat 1020  
 ggattcttca aatggaattt ctgaatacct ggatctatct tcccctggac cagtagaaat 1080  
 ttttaaccaca gtgacagaat cagttacagg tatttcaaga aatgtaagca ctaatgtggt 1140  
 cttcaagcaa catgattaca tcattgagtt ttttgattat actactgtct tgaagccatc 1200  
 tctcaacttc acagccactg tgaaggtaac tcgtgctgat ggcaaccaac tgactcttga 1260  
 agaaagaaga aataatgtag tcataacagt gacacagaga aactatactg agtactggag 1320  
 cggatctaac agtggaaatc agaaaatgga agctgttcag aaaataaatt atactgtccc 1380  
 ccaaagtgga acttttaaga ttgaattccc aatcctggag gattccagtg agctacagtt 1440  
 gaaggcctat ttccttggtg gtaaaagtag catggcagtt catagtctgt ttaagtctcc 1500

tagtaagaca tacatccaac taaaaacaag agatgaaaat ataaaggtgg gatcgcccttt	1560
tgagttggtg gttagtggca acaaacgatt gaaggagtta agctatatgg tagtatccag	1620
gggacagttg gtggctgtag gaaaacaaaa ttcaacaatg ttctctttta caccagaaaa	1680
ttcttggact caaaaagcct gtgtaattgt gtattatatt gaagatgatg gggaaattat	1740
aagtgatggt ctaaaaattc ctgttcagct tgtttttaaa aataagataa agctatatgt	1800
gagtaaagtg aaagctgaac catctgagaa agtctctctt aggatctctg tgacacagcc	1860
tgactccata gttgggattg tagctgttga caaaagtgtg aatctgatga atgcctctaa	1920
tgatattaca atggaaaatg tgggccatga gttggaactt tataacacag gatattattt	1980
aggcatgttc atgaattctt ttgcagtctt tcaggaatgt ggactctggg tattgacaga	2040
tgcaaacctc acgaaggatt atattgatgg tgtttatgac aatgcagaat atgctgagag	2100
gtttatggag gaaaatgaag gacatattgt agatattcat gacttttctt tgggtagcag	2160
tccacatgtc cgaaagcatt ttccagagac ttggatttgg ctagacacca acatgggttn	2220
caggatttac caagaatttg aagtaactgt acctgattct atcacttctt ggggtggctac	2280
tggttttgtg atctctgagg acctgggtct tggactaaca actactccag tggagctcca	2340
agccttccaa ccatttttca tttttttgaa tcttcctac tctgttatca gaggtgaaga	2400
atttgctttg gaaataacta tattcaatta ttgaaagat gccactgagg ttaaggtaat	2460
cattgagaaa agtgacaaat ttgatattct aatgacttca aatgaaataa atgccacagg	2520
ccaccagcag acccttctgg ttcccagtga ggatggggca actgttcttt ttcccatcag	2580
gccaacacat ctgggagaaa ttccatatcac agtcacagct ctttcacca ctgcttctga	2640
tgctgtcacc cagatgattt tagtaaaggc tgaaggaata gaaaaatcat attcacaatc	2700
catcttatta gacttgactg acaataggct acagagtacc ctgaaaactt tgagtttctc	2760
atttcctcct aatacagtga ctggcagtga aagagttcag atcactgcaa ttggagatgt	2820
tcttggctct tccatcaatg gcttagcctc attgattcgg atgccttatg gctgtggtga	2880
acagaacatg ataaattttg ctccaaatat ttacattttg gattatctga ctaaaaagaa	2940
acaactgaca gataatttga aagaaaaagc tctttcattt atgaggcaag gttaccagag	3000
agaacttctc tatcagaggg aagatggctc ttccagtgtc tttgggaatt atgacccttc	3060
tgggagcact tggttgtcag cttttgtttt aagatgtttc cttgaagccg atccttacat	3120
agatattgat cagaatgtgt tacacagaac atacacttgg cttaaaggac atcagaaatc	3180
caacggtgaa ttttgggatc caggaagagt gattcatagt gagcttcaag gtggcaataa	3240
aagtccagta acatttacag cctatattgt aacttctctc ctgggatata gaaagtatca	3300
gcctaacatt gatgtgcaag agtctatcca ttttttggag tctgaattca gtagaggaat	3360

ttcagacaat tatactctag cccttataac ttatgcattg tcatcagtg ggagtcctaa	3420
agcgaaggaa gctttgaata tgctgacttg gagagcagaa caagaagggtg gcatgcaatt	3480
ctgggtgtca tcagagtcca aactttctga ctctggcag ccacgctccc tggatattga	3540
agttgcagcc tatgcactgc tctcacactt cttacaattt cagacttctg agggaatccc	3600
aattatgagg tggctaagca ggcaaagaaa tagcttgggt ggttttgcat ctactcagga	3660
taccactgtg gctttaaagg ctctgtctga atttgcagcc ctaatgaata cagaaaggac	3720
aaatatccaa gtgaccgtga cggggcctag ctcaccaagt cctgtaaagt ttctgattga	3780
cacacacaac cgcttactcc ttcagacagc agagcttgct gtggtacagc caatggcagt	3840
taatatttcc gcaaatgggt ttggatttgc tatttgtcag ctcaatgttg tatataatgt	3900
gaaggcttct ggggtcttcta gaagacgaag atctatccaa aatcaagaag cctttgattt	3960
agatgttgct gtaaaagaaa ataaagatga tctcaatcat gtggatttga atgtgtgtac	4020
aagcttttctg ggcccgggtg ggagtggcat ggctcttatg gaagttaacc tattaagtgg	4080
ctttatgggtg ccttcagaag caatttctct gagcgagaca gtgaagaaag tggaatatga	4140
tcatggaaaa ctcaacctct atttagattc tgtaaatgaa acccagtttt gtgttaatat	4200
tcctgctgtg agaaacttta aagtttcaaa tacccaagat gcttcagtgt ccatagtgga	4260
ttactatgag ccaaggagac aggcggtgag aagttacaac tctgaagtga agctgtcctc	4320
ctgtgacctt tgcagtgatg tccagggtg ccgtccttgt gaggatggag cttcaggctc	4380
ccatcatcac tcttcagtca tttttatttt ctgtttcaag cttctgtact ttatggaact	4440
ttggctgtga tttattttta aaggactctg tgtaacacta acatttccag tagtcacatg	4500
tgattgtttt gttttcgtag aagaatactg cttctatttt gaaaaagag ttttttttct	4560
ttctatgggg ttgcagggat ggtgtacaac aggtcctagc atgtatagct gcatagattt	4620
cttcacctga tctttgtgtg gaagatcaga atgaatgcag ttgtgtgtct atattttccc	4680
ctcacaaaat cttttagaat ttttttgag gtgtttgttt tctccagaat aaaggtatta	4740
ctttagaaat aggtattctc ctcattttgt gaaagaaatg aacctagatt ctttaagcatt	4800
attacacatc catgtttgt taaagatgga tttccctggg aatgggagaa aacagccagc	4860
aggaggagct tcatctgttc ccttcccacc tccaacctag ccctactgcc caccacccc	4920
caaccacccc catgcccagt ggtctcagta gatacttctt aactggaaat tctttctttt	4980
cagaatctag gtggtgaatt ttttttaagt ggcacggtct ttttctgctt gaaatctgat	5040
cacacccccc agccattgcc ctccctctct ttttctctg tagagaaatg tgaggggcag	5100
tacatttact gtgcttttca caccatctca gaggttgagg agcatactga aaattgccct	5160
gggggggtgct ggggtgtgtg tctccttccc acatcctcag cccacacca gctctatttc	5220

- 13 -

```

aggggtgaga gtcagagagc actgcaatat gtgcttcgat ggatttcgat togaagatcc 5280
tagaccaggg agacactgtg agccagggat acaacaaaat actaggtaag tcaactgcaga 5340
ccgacctccc tgcagtttgg gaaagaagct gggtttgtgg agaatacagag catcttgaca 5400
tgactgctga cctaaagatc cctggcattg gccagggatc ctgtggaacc tcttctagtt 5460
caggggtgtg agcattagac tgccagttgt ctagtgacat ctgatgcttg ctgtgaactt 5520
ttaagatccc cgaatcctga gcacctcaat cttaattgc cctgtattcc gaagggtaat 5580
ataatttata tggatggaaa ttttaaagat gaatccccct ttttctttt cttctctctt 5640
ttctttcctt ctccctttct tctttgcctt ctaaatatac tgaaatgatt tagatatgtg 5700
tcaacaatta atgatctttt attcaatcta agaaatgggt tagtttttct ctttagctct 5760
atggcatttc actcaagtgg acaggggaaa aagtaattgc catgggctcc aaagaatttg 5820
ctttatgttt ttagctattt aaaaataaat ccatcaaaaa taaagtatgc aaatgtatct 5880
ttt 5883

```

```

<210> 4
<211> 1445
<212> PRT
<213> Homo sapiens

<220>
<221> MISC FEATURE
<222> (703)..(703)
<223> Xaa = Ser or Tyr

```

&lt;400&gt; 4

```

Met Gln Gly Pro Pro Leu Leu Thr Ala Ala His Leu Leu Cys Val Cys
1          5          10          15

```

```

Thr Ala Ala Leu Ala Val Ala Pro Gly Pro Arg Phe Leu Val Thr Ala
20          25          30

```

```

Pro Gly Ile Ile Arg Pro Gly Gly Asn Val Thr Ile Gly Val Glu Leu
35          40          45

```

```

Leu Glu His Cys Pro Ser Gln Val Thr Val Lys Ala Glu Leu Leu Lys
50          55          60

```

```

Thr Ala Ser Asn Leu Thr Val Ser Val Leu Glu Ala Glu Gly Val Phe
65          70          75          80

```

```

Glu Lys Gly Ser Phe Lys Thr Leu Thr Leu Pro Ser Leu Pro Leu Asn
85          90          95

```

- 14 -

Ser Ala Asp Glu Ile Tyr Glu Leu Arg Val Thr Gly Arg Thr Gln Asp  
 100 105 110

Glu Ile Leu Phe Ser Asn Ser Thr Arg Leu Ser Phe Glu Thr Lys Arg  
 115 120 125

Ile Ser Val Phe Ile Gln Thr Asp Lys Ala Leu Tyr Lys Pro Lys Gln  
 130 135 140

Glu Val Lys Phe Arg Ile Val Thr Leu Phe Ser Asp Phe Lys Pro Tyr  
 145 150 155 160

Lys Thr Ser Leu Asn Ile Leu Ile Lys Asp Pro Lys Ser Asn Leu Ile  
 165 170 175

Gln Gln Trp Leu Ser Gln Gln Ser Asp Leu Gly Val Ile Ser Lys Thr  
 180 185 190

Phe Gln Leu Ser Ser His Pro Ile Leu Gly Asp Trp Ser Ile Gln Val  
 195 200 205

Gln Val Asn Asp Gln Thr Tyr Tyr Gln Ser Phe Gln Val Ser Glu Tyr  
 210 215 220

Val Leu Pro Lys Phe Glu Val Thr Leu Gln Thr Pro Leu Tyr Cys Ser  
 225 230 235 240

Met Asn Ser Lys His Leu Asn Gly Thr Ile Thr Ala Lys Tyr Thr Tyr  
 245 250 255

Gly Lys Pro Val Lys Gly Asp Val Thr Leu Thr Phe Leu Pro Leu Ser  
 260 265 270

Phe Trp Gly Lys Lys Lys Asn Ile Thr Lys Thr Phe Lys Ile Asn Gly  
 275 280 285

Ser Ala Asn Phe Ser Phe Asn Asp Glu Glu Met Lys Asn Val Met Asp  
 290 295 300

Ser Ser Asn Gly Leu Ser Glu Tyr Leu Asp Leu Ser Ser Pro Gly Pro  
 305 310 315 320

Val Glu Ile Leu Thr Thr Val Thr Glu Ser Val Thr Gly Ile Ser Arg  
 325 330 335

Asn Val Ser Thr Asn Val Phe Phe Lys Gln His Asp Tyr Ile Ile Glu



340					345					350					
Phe	Phe	Asp	Tyr	Thr	Thr	Val	Leu	Lys	Pro	Ser	Leu	Asn	Phe	Thr	Ala
		355					360					365			
Thr	Val	Lys	Val	Thr	Arg	Ala	Asp	Gly	Asn	Gln	Leu	Thr	Leu	Glu	Glu
	370					375					380				
Arg	Arg	Asn	Asn	Val	Val	Ile	Thr	Val	Thr	Gln	Arg	Asn	Tyr	Thr	Glu
385					390					395					400
Tyr	Trp	Ser	Gly	Ser	Asn	Ser	Gly	Asn	Gln	Lys	Met	Glu	Ala	Val	Gln
				405					410					415	
Lys	Ile	Asn	Tyr	Thr	Val	Pro	Gln	Ser	Gly	Thr	Phe	Lys	Ile	Glu	Phe
			420					425					430		
Pro	Ile	Leu	Glu	Asp	Ser	Ser	Glu	Leu	Gln	Leu	Lys	Ala	Tyr	Phe	Leu
		435					440					445			
Gly	Ser	Lys	Ser	Ser	Met	Ala	Val	His	Ser	Leu	Phe	Lys	Ser	Pro	Ser
	450					455					460				
Lys	Thr	Tyr	Ile	Gln	Leu	Lys	Thr	Arg	Asp	Glu	Asn	Ile	Lys	Val	Gly
465					470					475					480
Ser	Pro	Phe	Glu	Leu	Val	Val	Ser	Gly	Asn	Lys	Arg	Leu	Lys	Glu	Leu
				485					490					495	
Ser	Tyr	Met	Val	Val	Ser	Arg	Gly	Gln	Leu	Val	Ala	Val	Gly	Lys	Gln
			500					505					510		
Asn	Ser	Thr	Met	Phe	Ser	Leu	Thr	Pro	Glu	Asn	Ser	Trp	Thr	Pro	Lys
		515					520					525			
Ala	Cys	Val	Ile	Val	Tyr	Tyr	Ile	Glu	Asp	Asp	Gly	Glu	Ile	Ile	Ser
	530					535					540				
Asp	Val	Leu	Lys	Ile	Pro	Val	Gln	Leu	Val	Phe	Lys	Asn	Lys	Ile	Lys
545					550					555					560
Leu	Tyr	Trp	Ser	Lys	Val	Lys	Ala	Glu	Pro	Ser	Glu	Lys	Val	Ser	Leu
				565					570					575	
Arg	Ile	Ser	Val	Thr	Gln	Pro	Asp	Ser	Ile	Val	Gly	Ile	Val	Ala	Val
			580					585					590		

- 16 -

Asp Lys Ser Val Asn Leu Met Asn Ala Ser Asn Asp Ile Thr Met Glu  
595 600 605

Asn Val Val His Glu Leu Glu Leu Tyr Asn Thr Gly Tyr Tyr Leu Gly  
610 615 620

Met Phe Met Asn Ser Phe Ala Val Phe Gln Glu Cys Gly Leu Trp Val  
625 630 635 640

Leu Thr Asp Ala Asn Leu Thr Lys Asp Tyr Ile Asp Gly Val Tyr Asp  
645 650 655

Asn Ala Glu Tyr Ala Glu Arg Phe Met Glu Glu Asn Glu Gly His Ile  
660 665 670

Val Asp Ile His Asp Phe Ser Leu Gly Ser Ser Pro His Val Arg Lys  
675 680 685

His Phe Pro Glu Thr Trp Ile Trp Leu Asp Thr Asn Met Gly Xaa Arg  
690 695 700

Ile Tyr Gln Glu Phe Glu Val Thr Val Pro Asp Ser Ile Thr Ser Trp  
705 710 715 720

Val Ala Thr Gly Phe Val Ile Ser Glu Asp Leu Gly Leu Gly Leu Thr  
725 730 735

Thr Thr Pro Val Glu Leu Gln Ala Phe Gln Pro Phe Phe Ile Phe Leu  
740 745 750

Asn Leu Pro Tyr Ser Val Ile Arg Gly Glu Glu Phe Ala Leu Glu Ile  
755 760 765

Thr Ile Phe Asn Tyr Leu Lys Asp Ala Thr Glu Val Lys Val Ile Ile  
770 775 780

Glu Lys Ser Asp Lys Phe Asp Ile Leu Met Thr Ser Asn Glu Ile Asn  
785 790 795 800

Ala Thr Gly His Gln Gln Thr Leu Leu Val Pro Ser Glu Asp Gly Ala  
805 810 815

Thr Val Leu Phe Pro Ile Arg Pro Thr His Leu Gly Glu Ile Pro Ile  
820 825 830

Thr Val Thr Ala Leu Ser Pro Thr Ala Ser Asp Ala Val Thr Gln Met

- 17 -

835	840	845
Ile Leu Val Lys Ala Glu Gly Ile Glu Lys Ser Tyr Ser Gln Ser Ile		
850	855	860
Leu Leu Asp Leu Thr Asp Asn Arg Leu Gln Ser Thr Leu Lys Thr Leu		
865	870	875 880
Ser Phe Ser Phe Pro Pro Asn Thr Val Thr Gly Ser Glu Arg Val Gln		
	885	890 895
Ile Thr Ala Ile Gly Asp Val Leu Gly Pro Ser Ile Asn Gly Leu Ala		
	900	905 910
Ser Leu Ile Arg Met Pro Tyr Gly Cys Gly Glu Gln Asn Met Ile Asn		
	915	920 925
Phe Ala Pro Asn Ile Tyr Ile Leu Asp Tyr Leu Thr Lys Lys Lys Gln		
	930	935 940
Leu Thr Asp Asn Leu Lys Glu Lys Ala Leu Ser Phe Met Arg Gln Gly		
945	950	955 960
Tyr Gln Arg Glu Leu Leu Tyr Gln Arg Glu Asp Gly Ser Phe Ser Ala		
	965	970 975
Phe Gly Asn Tyr Asp Pro Ser Gly Ser Thr Trp Leu Ser Ala Phe Val		
	980	985 990
Leu Arg Cys Phe Leu Glu Ala Asp Pro Tyr Ile Asp Ile Asp Gln Asn		
	995	1000 1005
Val Leu His Arg Thr Tyr Thr Trp Leu Lys Gly His Gln Lys Ser		
1010	1015	1020
Asn Gly Glu Phe Trp Asp Pro Gly Arg Val Ile His Ser Glu Leu		
1025	1030	1035
Gln Gly Gly Asn Lys Ser Pro Val Thr Leu Thr Ala Tyr Ile Val		
1040	1045	1050
Thr Ser Leu Leu Gly Tyr Arg Lys Tyr Gln Pro Asn Ile Asp Val		
1055	1060	1065
Gln Glu Ser Ile His Phe Leu Glu Ser Glu Phe Ser Arg Gly Ile		
1070	1075	1080

- 18 -

Ser	Asp	Asn	Tyr	Thr	Leu	Ala	Leu	Ile	Thr	Tyr	Ala	Leu	Ser	Ser
1085						1090					1095			
Val	Gly	Ser	Pro	Lys	Ala	Lys	Glu	Ala	Leu	Asn	Met	Leu	Thr	Trp
1100						1105					1110			
Arg	Ala	Glu	Gln	Glu	Gly	Gly	Met	Gln	Phe	Trp	Val	Ser	Ser	Glu
1115						1120					1125			
Ser	Lys	Leu	Ser	Asp	Ser	Trp	Gln	Pro	Arg	Ser	Leu	Asp	Ile	Glu
1130						1135					1140			
Val	Ala	Ala	Tyr	Ala	Leu	Leu	Ser	His	Phe	Leu	Gln	Phe	Gln	Thr
1145						1150					1155			
Ser	Glu	Gly	Ile	Pro	Ile	Met	Arg	Trp	Leu	Ser	Arg	Gln	Arg	Asn
1160						1165					1170			
Ser	Leu	Gly	Gly	Phe	Ala	Ser	Thr	Gln	Asp	Thr	Thr	Val	Ala	Leu
1175						1180					1185			
Lys	Ala	Leu	Ser	Glu	Phe	Ala	Ala	Leu	Met	Asn	Thr	Glu	Arg	Thr
1190						1195					1200			
Asn	Ile	Gln	Val	Thr	Val	Thr	Gly	Pro	Ser	Ser	Pro	Ser	Pro	Val
1205						1210					1215			
Lys	Phe	Leu	Ile	Asp	Thr	His	Asn	Arg	Leu	Leu	Leu	Gln	Thr	Ala
1220						1225					1230			
Glu	Leu	Ala	Val	Val	Gln	Pro	Met	Ala	Val	Asn	Ile	Ser	Ala	Asn
1235						1240					1245			
Gly	Phe	Gly	Phe	Ala	Ile	Cys	Gln	Leu	Asn	Val	Val	Tyr	Asn	Val
1250						1255					1260			
Lys	Ala	Ser	Gly	Ser	Ser	Arg	Arg	Arg	Arg	Ser	Ile	Gln	Asn	Gln
1265						1270					1275			
Glu	Ala	Phe	Asp	Leu	Asp	Val	Ala	Val	Lys	Glu	Asn	Lys	Asp	Asp
1280						1285					1290			
Leu	Asn	His	Val	Asp	Leu	Asn	Val	Cys	Thr	Ser	Phe	Ser	Gly	Pro
1295						1300					1305			
Gly	Arg	Ser	Gly	Met	Ala	Leu	Met	Glu	Val	Asn	Leu	Leu	Ser	Gly

- 19 -

1310	1315	1320
Phe Met Val Pro Ser Glu Ala Ile Ser Leu Ser Glu Thr Val Lys		
1325	1330	1335
Lys Val Glu Tyr Asp His Gly Lys Leu Asn Leu Tyr Leu Asp Ser		
1340	1345	1350
Val Asn Glu Thr Gln Phe Cys Val Asn Ile Pro Ala Val Arg Asn		
1355	1360	1365
Phe Lys Val Ser Asn Thr Gln Asp Ala Ser Val Ser Ile Val Asp		
1370	1375	1380
Tyr Tyr Glu Pro Arg Arg Gln Ala Val Arg Ser Tyr Asn Ser Glu		
1385	1390	1395
Val Lys Leu Ser Ser Cys Asp Leu Cys Ser Asp Val Gln Gly Cys		
1400	1405	1410
Arg Pro Cys Glu Asp Gly Ala Ser Gly Ser His His His Ser Ser		
1415	1420	1425
Val Ile Phe Ile Phe Cys Phe Lys Leu Leu Tyr Phe Met Glu Leu		
1430	1435	1440
Trp Leu		
1445		

<210> 5  
 <211> 4146  
 <212> DNA  
 <213> Homo sapiens  
 <220>  
 <221> misc feature  
 <222> (2045)..(2045)  
 <223> N = A or C

<400> 5	
gtggctcccg ggccctcggtt tctggtgaca gccccagggg tcatcaggcc cggaggaaat	60
gtgactattg ggggtggagct tctggaacac tgcccttcac aggtgactgt gaaggcggag	120
ctgctcaaga cagcatcaaa cctcactgtc tctgtcctgg aagcagaagg agtctttgaa	180
aaaggctctt ttaagacact tactcttcca tcactacctc tgaacagtgc agatgagatt	240
tatgagctac gtgtaaccgg acgtacccag gatgagattt tattctctaa tagtaccgcg	300
ttatcatttg agaccaagag aatatctgtc ttcatcctaaa cagacaaggc cttatacaag	360

ccaaagcaag aagtgaagtt togcattggt acactcttct cagattttta gccttacaaa	420
acctctttta acattctcat taaggacccc aaatcaaatt tgatccaaca gtgggtgtca	480
caacaaagtg atcttggagt catttccaaa acttttcagc tatcttccca tocaatactt	540
ggtgactggt ctattcaagt tcaagtgaat gaccagacat actatcaatc atttcagggt	600
tcagaatatg tattacccaa atttgaagtg actttgcaga caccattata ttgttctatg	660
aattctaagc atttaaatgg taccatcacg gcaaagtata catatgggaa gccagtga	720
ggagacgtaa cgcttacatt ttaccttta tccttttggg gaaagaagaa aaatattaca	780
aaaacattta agataaatgg atctgcaaac ttctctttta atgatgaaga gatgaaaaat	840
gtaatggatt cttcaaatgg actttctgaa tacctggatc tatcttcccc tggaccagta	900
gaaattttta ccacagtgc agaatcagtt acaggatatt caagaaatgt aagcactaat	960
gtgttcttca agcaacatga ttacatcatt gagttttttg attatactac tgtcttgaag	1020
ccatctctca acttcacagc cactgtgaag gtaactcgtg ctgatggcaa ccaactgact	1080
cttgaagaaa gaagaaataa tgtagtata acagtgcac agagaaacta tactgagtac	1140
tggagcggat ctaacagtgg aaatcagaaa atggaagctg ttcagaaaat aaattatact	1200
gtccccaaa gtggaacttt taagattgaa ttcccaatcc tggaggattc cagtgcgta	1260
cagttgaagg cctatttctt tggtagtaaa agtagcatgg cagttcatag tctgtttaag	1320
tctcctagta agacatacat ccaactaaaa acaagagatg aaaatataaa ggtgggatcg	1380
ccttttgagt tgggtggttag tggcaacaaa cgattgaagg agttaagcta tatggtagta	1440
tccaggggac agttggtggc tgtaggaaaa caaaattcaa caatgttctc tttaacacca	1500
gaaaattctt ggactccaaa agcctgtgta attgtgtatt atattgaaga tgatggggaa	1560
attataagtg atgttctaaa aattcctgtt cagcttggtt ttaaaaataa gataaagcta	1620
tattggagta aagtgaagc tgaaccatct gagaaagtct ctcttaggat ctctgtgaca	1680
cagcctgact ccatagttgg gattgtagct gttgacaaaa gtgtgaatct gatgaatgcc	1740
tctaatagata ttacaatgga aatgtggtc catgagttgg aactttataa cacaggatat	1800
tatttaggca tgttcatgaa ttcttttgca gtctttcagg aatgtggact ctgggtattg	1860
acagatgcaa acctcacgaa ggattatatt gatggtggtt atgacaatgc agaatatgct	1920
gagaggttta tggaggaaaa tgaaggacat attgtagata ttcatgactt ttctttgggt	1980
agcagtcac atgtccgaaa gcattttcca gagacttga tttggctaga caccaacatg	2040
ggttncagga ttaccaaga atttgaagta actgtacctg attctatcac ttcttgggtg	2100
gctactggtt ttgtgatctc tgaggacctg ggtcttggac taacaactac tccagtggag	2160
ctccaagcct tccaaccatt ttcatTTTT ttgaatcttc cctactctgt tatcagaggt	2220

gaagaatttg	ctttggaaat	aactatat	aattatttga	aagatgccac	tgagggttaag	2280
gtaatcattg	agaaaagtga	caaatttgat	attctaata	cttcaaata	aataaatgcc	2340
acaggccacc	agcagaccct	tctggttccc	agtgaggatg	gggcaactgt	tctttttccc	2400
atcaggccaa	cacatctggg	agaaattcct	atcacagtca	cagctctttc	acctactgct	2460
tctgatgctg	tcaccagat	gattttagta	aaggctgaag	gaatagaaaa	atcatattca	2520
caatccatct	tattagactt	gactgacaat	aggctacaga	gtaccctgaa	aactttgagt	2580
ttctcatttc	ctcctaatac	agtgactggc	agtgaaagag	ttcagatcac	tgcaattgga	2640
gatgttcttg	gtccttccat	caatggctta	gcctcattga	ttcggatgcc	ttatggctgt	2700
ggtgaacaga	acatgataaa	ttttgctcca	aatattttaca	ttttggatta	tctgactaaa	2760
aagaaacaac	tgacagataa	tttgaaagaa	aaagctcttt	catttatgag	gcaagggttac	2820
cagagagAAC	ttctctatca	gagggaagat	ggctctttca	gtgcttttgg	gaattatgac	2880
ccttctggga	gcacttggtt	gtcagctttt	gttttaagat	gtttccttga	agccgatcct	2940
tacatagata	ttgatcagaa	tgtgttacac	agaacataca	cttggcttaa	aggacatcag	3000
aaatccaacg	gtgaattttg	ggatccagga	agagtgattc	atagtgagct	tcaagggtggc	3060
aataaaaagtc	cagtaacact	tacagcctat	attgtaactt	ctctcctggg	atatagaaaag	3120
tatcagccta	acattgatgt	gcaagagtct	atccattttt	tggagtctga	attcagtaga	3180
ggaatttcag	acaattatac	tctagccctt	ataacttatg	cattgtcatc	agtggggagt	3240
cctaaagcga	aggaagcttt	gaatatgctg	acttgagag	cagaacaaga	aggtggcatg	3300
caattctggg	tgatcatcaga	gtccaaactt	tctgactcct	ggcagccacg	ctccctggat	3360
attgaagttg	cagcctatgc	actgctctca	cacttcttac	aatttcagac	ttctgaggga	3420
atcccaatta	tgagggtggct	aagcaggcaa	agaaatagct	tgggtggttt	tgcatctact	3480
caggatacca	ctgtggcttt	aaaggctctg	totgaatttg	cagccctaata	gaatacagaa	3540
aggacaaata	tccaagtgc	cgtgacggg	cctagctcac	caagtcctct	tgctgtggta	3600
cagccaacgg	cagttaatat	ttccgcaaat	ggttttgat	ttgctatttg	tcagctcaat	3660
gttgatatata	atgtgaaggc	ttctgggtct	tctagaagac	gaagatctat	ccaaaatcaa	3720
gaagcctttg	atttagatgt	tgctgtaaaa	gaaaataaag	atgatctcaa	tcagtgtgat	3780
ttgaatgtgt	gtacaagctt	ttcgggccc	ggtaggagt	gcatggctct	tatggaagtt	3840
aacctattaa	gtggctttat	ggtgccttca	gaagcaattt	ctctgagcga	gacagtgaag	3900
aaagtggaat	atgatcatgg	aaaactcaac	ctctattttag	attctgtaaa	tgaaaccacg	3960
ttttgtgtta	atattcctgc	tgtgagaaac	tttaaagttt	caaataccca	agatgcttca	4020
gtgtccatag	tggattacta	tgagccaagg	agacaggcgg	tgagaagtta	caactctgaa	4080

- 22 -

gtgaagctgt cctcctgtga cctttgcagt gatgtccagg gctgccgtcc ttgtgaggat 4140  
 ggagct 4146

<210> 6  
 <211> 1382  
 <212> PRT  
 <213> homo sapiens  
 <220>  
 <221> MISC\_FEATURE  
 <222> (682)..(682)  
 <223> Xaa = Ser or Tyr

<400> 6

Val Ala Pro Gly Pro Arg Phe Leu Val Thr Ala Pro Gly Ile Ile Arg  
 1 5 10 15

Pro Gly Gly Asn Val Thr Ile Gly Val Glu Leu Leu Glu His Cys Pro  
 20 25 30

Ser Gln Val Thr Val Lys Ala Glu Leu Leu Lys Thr Ala Ser Asn Leu  
 35 40 45

Thr Val Ser Val Leu Glu Ala Glu Gly Val Phe Glu Lys Gly Ser Phe  
 50 55 60

Lys Thr Leu Thr Leu Pro Ser Leu Pro Leu Asn Ser Ala Asp Glu Ile  
 65 70 75 80

Tyr Glu Leu Arg Val Thr Gly Arg Thr Gln Asp Glu Ile Leu Phe Ser  
 85 90 95

Asn Ser Thr Arg Leu Ser Phe Glu Thr Lys Arg Ile Ser Val Phe Ile  
 100 105 110

Gln Thr Asp Lys Ala Leu Tyr Lys Pro Lys Gln Glu Val Lys Phe Arg  
 115 120 125

Ile Val Thr Leu Phe Ser Asp Phe Lys Pro Tyr Lys Thr Ser Leu Asn  
 130 135 140

Ile Leu Ile Lys Asp Pro Lys Ser Asn Leu Ile Gln Gln Trp Leu Ser  
 145 150 155 160

Gln Gln Ser Asp Leu Gly Val Ile Ser Lys Thr Phe Gln Leu Ser Ser  
 165 170 175



- 23 -

His Pro Ile Leu Gly Asp Trp Ser Ile Gln Val Gln Val Asn Asp Gln  
180 185 190

Thr Tyr Tyr Gln Ser Phe Gln Val Ser Glu Tyr Val Leu Pro Lys Phe  
195 200 205

Glu Val Thr Leu Gln Thr Pro Leu Tyr Cys Ser Met Asn Ser Lys His  
210 215 220

Leu Asn Gly Thr Ile Thr Ala Lys Tyr Thr Tyr Gly Lys Pro Val Lys  
225 230 235 240

Gly Asp Val Thr Leu Thr Phe Leu Pro Leu Ser Phe Trp Gly Lys Lys  
245 250 255

Lys Asn Ile Thr Lys Thr Phe Lys Ile Asn Gly Ser Ala Asn Phe Ser  
260 265 270

Phe Asn Asp Glu Glu Met Lys Asn Val Met Asp Ser Ser Asn Gly Leu  
275 280 285

Ser Glu Tyr Leu Asp Leu Ser Ser Pro Gly Pro Val Glu Ile Leu Thr  
290 295 300

Thr Val Thr Glu Ser Val Thr Gly Ile Ser Arg Asn Val Ser Thr Asn  
305 310 315 320

Val Phe Phe Lys Gln His Asp Tyr Ile Ile Glu Phe Phe Asp Tyr Thr  
325 330 335

Thr Val Leu Lys Pro Ser Leu Asn Phe Thr Ala Thr Val Lys Val Thr  
340 345 350

Arg Ala Asp Gly Asn Gln Leu Thr Leu Glu Glu Arg Arg Asn Asn Val  
355 360 365

Val Ile Thr Val Thr Gln Arg Asn Tyr Thr Glu Tyr Trp Ser Gly Ser  
370 375 380

Asn Ser Gly Asn Gln Lys Met Glu Ala Val Gln Lys Ile Asn Tyr Thr  
385 390 395 400

Val Pro Gln Ser Gly Thr Phe Lys Ile Glu Phe Pro Ile Leu Glu Asp  
405 410 415

Ser Ser Glu Leu Gln Leu Lys Ala Tyr Phe Leu Gly Ser Lys Ser Ser

- 24 -

420	425	430
Met Ala Val His Ser Leu Phe Lys Ser Pro Ser Lys Thr Tyr Ile Gln 435 440 445		
Leu Lys Thr Arg Asp Glu Asn Ile Lys Val Gly Ser Pro Phe Glu Leu 450 455 460		
Val Val Ser Gly Asn Lys Arg Leu Lys Glu Leu Ser Tyr Met Val Val 465 470 475 480		
Ser Arg Gly Gln Leu Val Ala Val Gly Lys Gln Asn Ser Thr Met Phe 485 490 495		
Ser Leu Thr Pro Glu Asn Ser Trp Thr Pro Lys Ala Cys Val Ile Val 500 505 510		
Tyr Tyr Ile Glu Asp Asp Gly Glu Ile Ile Ser Asp Val Leu Lys Ile 515 520 525		
Pro Val Gln Leu Val Phe Lys Asn Lys Ile Lys Leu Tyr Trp Ser Lys 530 535 540		
Val Lys Ala Glu Pro Ser Glu Lys Val Ser Leu Arg Ile Ser Val Thr 545 550 555 560		
Gln Pro Asp Ser Ile Val Gly Ile Val Ala Val Asp Lys Ser Val Asn 565 570 575		
Leu Met Asn Ala Ser Asn Asp Ile Thr Met Glu Asn Val Val His Glu 580 585 590		
Leu Glu Leu Tyr Asn Thr Gly Tyr Tyr Leu Gly Met Phe Met Asn Ser 595 600 605		
Phe Ala Val Phe Gln Glu Cys Gly Leu Trp Val Leu Thr Asp Ala Asn 610 615 620		
Leu Thr Lys Asp Tyr Ile Asp Gly Val Tyr Asp Asn Ala Glu Tyr Ala 625 630 635 640		
Glu Arg Phe Met Glu Glu Asn Glu Gly His Ile Val Asp Ile His Asp 645 650 655		
Phe Ser Leu Gly Ser Ser Pro His Val Arg Lys His Phe Pro Glu Thr 660 665 670		

- 25 -

Trp Ile Trp Leu Asp Thr Asn Met Gly Xaa Arg Ile Tyr Gln Glu Phe  
 675 685

Glu Val Thr Val Pro Asp Ser Ile Thr Ser Trp Val Ala Thr Gly Phe  
 690 700

Val Ile Ser Glu Asp Leu Gly Leu Gly Leu Thr Thr Thr Pro Val Glu  
 705 710 715 720

Leu Gln Ala Phe Gln Pro Phe Phe Ile Phe Leu Asn Leu Pro Tyr Ser  
 725 730 735

Val Ile Arg Gly Glu Glu Phe Ala Leu Glu Ile Thr Ile Phe Asn Tyr  
 740 745 750

Leu Lys Asp Ala Thr Glu Val Lys Val Ile Ile Glu Lys Ser Asp Lys  
 755 760 765

Phe Asp Ile Leu Met Thr Ser Asn Glu Ile Asn Ala Thr Gly His Gln  
 770 775 780

Gln Thr Leu Leu Val Pro Ser Glu Asp Gly Ala Thr Val Leu Phe Pro  
 785 790 795 800

Ile Arg Pro Thr His Leu Gly Glu Ile Pro Ile Thr Val Thr Ala Leu  
 805 810 815

Ser Pro Thr Ala Ser Asp Ala Val Thr Gln Met Ile Leu Val Lys Ala  
 820 825 830

Glu Gly Ile Glu Lys Ser Tyr Ser Gln Ser Ile Leu Leu Asp Leu Thr  
 835 840 845

Asp Asn Arg Leu Gln Ser Thr Leu Lys Thr Leu Ser Phe Ser Phe Pro  
 850 855 860

Pro Asn Thr Val Thr Gly Ser Glu Arg Val Gln Ile Thr Ala Ile Gly  
 865 870 875 880

Asp Val Leu Gly Pro Ser Ile Asn Gly Leu Ala Ser Leu Ile Arg Met  
 885 890 895

Pro Tyr Gly Cys Gly Glu Gln Asn Met Ile Asn Phe Ala Pro Asn Ile  
 900 905 910

Tyr Ile Leu Asp Tyr Leu Thr Lys Lys Lys Gln Leu Thr Asp Asn Leu

915	920	925
Lys Glu Lys Ala Leu Ser Phe Met Arg Gln Gly Tyr Gln Arg Glu Leu		
930	935	940
Leu Tyr Gln Arg Glu Asp Gly Ser Phe Ser Ala Phe Gly Asn Tyr Asp		
945	950	955 960
Pro Ser Gly Ser Thr Trp Leu Ser Ala Phe Val Leu Arg Cys Phe Leu		
	965	970 975
Glu Ala Asp Pro Tyr Ile Asp Ile Asp Gln Asn Val Leu His Arg Thr		
	980	985 990
Tyr Thr Trp Leu Lys Gly His Gln Lys Ser Asn Gly Glu Phe Trp Asp		
	995	1000 1005
Pro Gly Arg Val Ile His Ser Glu Leu Gln Gly Gly Asn Lys Ser		
1010	1015	1020
Pro Val Thr Leu Thr Ala Tyr Ile Val Thr Ser Leu Leu Gly Tyr		
1025	1030	1035
Arg Lys Tyr Gln Pro Asn Ile Asp Val Gln Glu Ser Ile His Phe		
1040	1045	1050
Leu Glu Ser Glu Phe Ser Arg Gly Ile Ser Asp Asn Tyr Thr Leu		
1055	1060	1065
Ala Leu Ile Thr Tyr Ala Leu Ser Ser Val Gly Ser Pro Lys Ala		
1070	1075	1080
Lys Glu Ala Leu Asn Met Leu Thr Trp Arg Ala Glu Gln Glu Gly		
1085	1090	1095
Gly Met Gln Phe Trp Val Ser Ser Glu Ser Lys Leu Ser Asp Ser		
1100	1105	1110
Trp Gln Pro Arg Ser Leu Asp Ile Glu Val Ala Ala Tyr Ala Leu		
1115	1120	1125
Leu Ser His Phe Leu Gln Phe Gln Thr Ser Glu Gly Ile Pro Ile		
1130	1135	1140
Met Arg Trp Leu Ser Arg Gln Arg Asn Ser Leu Gly Gly Phe Ala		
1145	1150	1155

- 27 -

Ser Thr	Gln Asp Thr Thr	Val	Ala Leu Lys Ala	Leu	Ser Glu Phe
1160		1165		1170	
Ala Ala	Leu Met Asn Thr	Glu	Arg Thr Asn Ile	Gln	Val Thr Val
1175		1180		1185	
Thr Gly	Pro Ser Ser Pro	Ser	Pro Leu Ala Val	Val	Gln Pro Thr
1190		1195		1200	
Ala Val	Asn Ile Ser Ala	Asn	Gly Phe Gly Phe	Ala	Ile Cys Gln
1205		1210		1215	
Leu Asn	Val Val Tyr Asn	Val	Lys Ala Ser Gly	Ser	Ser Arg Arg
1220		1225		1230	
Arg Arg	Ser Ile Gln Asn	Gln	Glu Ala Phe Asp	Leu	Asp Val Ala
1235		1240		1245	
Val Lys	Glu Asn Lys Asp	Asp	Leu Asn His Val	Asp	Leu Asn Val
1250		1255		1260	
Cys Thr	Ser Phe Ser Gly	Pro	Gly Arg Ser Gly	Met	Ala Leu Met
1265		1270		1275	
Glu Val	Asn Leu Leu Ser	Gly	Phe Met Val Pro	Ser	Glu Ala Ile
1280		1285		1290	
Ser Leu	Ser Glu Thr Val	Lys	Lys Val Glu Tyr	Asp	His Gly Lys
1295		1300		1305	
Leu Asn	Leu Tyr Leu Asp	Ser	Val Asn Glu Thr	Gln	Phe Cys Val
1310		1315		1320	
Asn Ile	Pro Ala Val Arg	Asn	Phe Lys Val Ser	Asn	Thr Gln Asp
1325		1330		1335	
Ala Ser	Val Ser Ile Val	Asp	Tyr Tyr Glu Pro	Arg	Arg Gln Ala
1340		1345		1350	
Val Arg	Ser Tyr Asn Ser	Glu	Val Lys Leu Ser	Ser	Cys Asp Leu
1355		1360		1365	
Cys Ser	Asp Val Gln Gly	Cys	Arg Pro Cys Glu	Asp	Gly Ala
1370		1375		1380	

&lt;210&gt; 7

<211> 4197  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <222> (2045)..(2045)  
 <223> N = A or C

<400> 7  
 gtggctcccg ggcctcggtt tctggtgaca gccccagggg tcatcaggcc cggaggaaat 60  
 gtgactattg ggggtggagct tctggaacac tgcccttcac aggtgactgt gaaggcggag 120  
 ctgctcaaga cagcatcaaa cctcactgtc tctgtcctgg aagcagaagg agtctttgaa 180  
 aaaggctctt ttaagacact tactcttcca tcactacctc tgaacagtgc agatgagatt 240  
 tatgagctac gtgtaaccgg acgtaccag gatgagattt tattctctaa tagtaccgcg 300  
 ttatcatttg agaccaagag aatatctgtc ttcatcaca cagacaaggc cttatacaag 360  
 ccaaagcaag aagtgaagtt tcgcattggt acactcttct cagattttta gccttacaaa 420  
 acctctttta acattctcat taaggacccc aatcaaatt tgatccaaca gtgggtgtca 480  
 caacaaagt atcttgaggt catttccaaa acttttcagc tatcttccca tccaataact 540  
 ggtgactggt ctattcaagt tcaagtgaat gaccagacat attatcaatc atttcagggt 600  
 tcagaatatg tattaccaaa atttgaagt actttgcaga caccattata ttgttctatg 660  
 aattctaagc atttaaatgg taccatcacg gcaaagtata catatgggaa gccagtga 720  
 ggagacgtaa cgcttacatt ttaccttta tccttttggg gaaagaagaa aaatattaca 780  
 aaaacattta agataaatgg atctgcaaac ttctctttta atgatgaaga gatgaaaaat 840  
 gtaatggatt cttcaaagt actttctgaa tacctggatc tatcttcccc tggaccagta 900  
 gaaattttta ccacagtgc agaatcagtt acaggtattt caagaaatgt aagcactaat 960  
 gtgttcttca agcaacatga ttacatcatt gagttttttg attatactac tgtcttgaag 1020  
 ccatctctca acttcacagc cactgtgaag gtaactcgtg ctgatggcaa ccaactgact 1080  
 cttgaagaaa gaagaaataa tgtagtcata acagtgcac agagaaacta tactgagtac 1140  
 tggagcggat ctaacagtgg aaatcagaaa atggaagctg tcagaaaaat aaattatact 1200  
 gtccccaaa gtggaacttt taagattgaa ttcccaatcc tggaggattc cagtgaagta 1260  
 cagttgaagg cctatttcct tggtagtaaa agtagcatgg cagttcatag tctgtttaag 1320  
 tctcctagta agacatacat ccaactaaaa acaagagatg aaaatataaa ggtgggatcg 1380  
 ccttttgagt tgggtggttag tggcaacaaa cgattgaagg agttaagcta tatggtagta 1440  
 tccaggggac agttggtggc tgtaggaaaa caaaattcaa caatgttctc tttaacacca 1500  
 gaaaattctt ggactccaaa agcctgtgta attgtgtatt atattgaaga tgatggggaa 1560

attataagtg atgttctaaa aattcctggt cagcttggtt ttaaaaataa gataaagcta	1620
tattggagta aagtgaaagc tgaaccatct gagaaagtct ctcttaggat ctctgtgaca	1680
cagcctgact ccatagttgg gattgtagct gttgacaaaa gtgtgaatct gatgaatgcc	1740
tctaagtata ttacaatgga aaatgtggtc catgagttgg aactttataa cacaggatat	1800
tatttaggca tgttcatgaa ttcttttgca gtctttcagg aatgtggact ctgggtattg	1860
acagatgcaa acctcacgaa ggattatatt gatggtgttt atgacaatgc agaatatgct	1920
gagaggttta tggaggaaaa tgaaggacat attgtagata ttcattgactt ttctttgggt	1980
agcagtcac atgtccgaaa gcattttcca gagacttggga tttggctaga caccaacatg	2040
ggttncagga ttaccaaga atttgaagta actgtacctg attctatcac ttcttgggtg	2100
gctactgggt ttgtgatctc tgaggacctg ggtcttggac taacaactac tccagtggag	2160
ctccaagcct tccaaccatt ttctattttt ttgaatcttc cctactctgt tatcagaggt	2220
gaagaatttg ctttggaaat aactatattc aattatttga aagatgccac tgaggttaag	2280
gtaatcattg agaaaagtga caaatttgat attctaataa cttcaaataa aataaatgcc	2340
acaggccacc agcagaccct tctggttccc agtgaggatg gggcaactgt tctttttccc	2400
atcaggccaa cacatctggg agaaattcct atcacagtca cagctctttc acccaactgt	2460
tctgatgctg tcaccagat gatttttagta aaggctgaag gaatagaaaa atcatattca	2520
caatccatct tattagactt gactgacaat aggctacaga gtacctgaa aactttgagt	2580
ttctcatttc ctctaatac agtgactggc agtgaaagag ttcagatcac tgcaattgga	2640
gatgttcttg gtccctccat caatggctta gcctcattga ttoggatgcc ttatggctgt	2700
ggtgaacaga acatgataaa ttttgctcca aatatttaca ttttggatta tctgactaaa	2760
aagaaacaac tgacagataa tttgaagaa aaagctcttt catttatgag gcaagggttac	2820
cagagagaac ttctctatca gagggagat ggctctttca gtgcttttgga gaattatgac	2880
ccttctggga gcacttgggt gtcagctttt gttttaagat gtttccttga agccgatcct	2940
tacatagata ttgatcagaa tgtgttacac agaacataca cttggcttaa aggacatcag	3000
aatccaacg gtgaattttg ggatccagga agagtgattc atagtgagct tcaagggtggc	3060
aataaaagtc cagtaaacct tacagcctat attgtaactt ctctcctggg atatagaaaag	3120
tatcagccta acattgatgt gcaagagtct atccattttt tggagtctga attcagtaga	3180
ggaatttcag acaattatac tctagccctt ataacttatg cattgtcatc agtggggagt	3240
cctaaagcga aggaagcttt gaatatgctg acttggagag cagaacaaga aggtggcatg	3300
caattctggg tgtcatcaga gtccaaactt tctgactcct ggcagccacg ctccctggat	3360
attgaagttg cagcctatgc actgctctca cacttcttac aatttcagac ttctgagga	3420

```

atcccaatta tgagggtggct aagcaggcaa agaaatagct tgggtgggtt tgcattact 3480
caggatacca ctgtggcttt aaaggctctg tctgaatttg cagccctaata gaatacagaa 3540
aggacaaata tccaagtgaac cgtgacgggg cctagctcac caagtcctgt aaagtttctg 3600
attgacacac acaaccgctt actccttcag acagcagagc ttgctgtggg acagccaatg 3660
gcagttaata tttccgcaaa tggttttgga tttgctatgt gtcagctcaa tgttgtatat 3720
aatgtgaagg cttctgggtc ttctagaaga cgaagatcta tccaaaatca agaagccttt 3780
gatttagatg ttgctgtaaa agaaaataaa gatgatctca atcatgtgga tttgaatgtg 3840
tgtacaagct tttcggggccc gggtaggagt ggcattggctc ttatggaagt taacctatta 3900
agtggcttta tggcgccttc agaagcaatt tctctgagcg agacagtga gaaagtggaa 3960
tatgatcatg gaaaactcaa cctctattta gattctgtaa atgaaacca gttttgtgtt 4020
aatattcctg ctgtgagaaa ctttaaagtt tcaaataccc aagatgcttc agtgtccata 4080
gtggattact atgagccaag gagacaggcg gtgagaagtt acaactctga agtgaagctg 4140
tcctcctgtg acctttgcag tgatgtccag ggctgccgctc cttgtgagga tggagct 4197

```

```

<210> 8
<211> 1399
<212> PRT
<213> Homo sapiens

<220>
<221> MISC_FEATURE
<222> (682)..(682)
<223> Xaa = Ser or Tyr

```

```

<400> 8

```

```

Val Ala Pro Gly Pro Arg Phe Leu Val Thr Ala Pro Gly Ile Ile Arg
1      5      10      15

```

```

Pro Gly Gly Asn Val Thr Ile Gly Val Glu Leu Leu Glu His Cys Pro
20      25      30

```

```

Ser Gln Val Thr Val Lys Ala Glu Leu Leu Lys Thr Ala Ser Asn Leu
35      40      45

```

```

Thr Val Ser Val Leu Glu Ala Glu Gly Val Phe Glu Lys Gly Ser Phe
50      55      60

```

```

Lys Thr Leu Thr Leu Pro Ser Leu Pro Leu Asn Ser Ala Asp Glu Ile
65      70      75      80

```

```

Tyr Glu Leu Arg Val Thr Gly Arg Thr Gln Asp Glu Ile Leu Phe Ser

```



- 31 -

					85						90					95
Asn	Ser	Thr	Arg	Leu	Ser	Phe	Glu	Thr	Lys	Arg	Ile	Ser	Val	Phe	Ile	
					100						105				110	
Gln	Thr	Asp	Lys	Ala	Leu	Tyr	Lys	Pro	Lys	Gln	Glu	Val	Lys	Phe	Arg	
		115					120					125				
Ile	Val	Thr	Leu	Phe	Ser	Asp	Phe	Lys	Pro	Tyr	Lys	Thr	Ser	Leu	Asn	
	130					135					140					
Ile	Leu	Ile	Lys	Asp	Pro	Lys	Ser	Asn	Leu	Ile	Gln	Gln	Trp	Leu	Ser	
145					150					155					160	
Gln	Gln	Ser	Asp	Leu	Gly	Val	Ile	Ser	Lys	Thr	Phe	Gln	Leu	Ser	Ser	
				165					170					175		
His	Pro	Ile	Leu	Gly	Asp	Trp	Ser	Ile	Gln	Val	Gln	Val	Asn	Asp	Gln	
			180					185					190			
Thr	Tyr	Tyr	Gln	Ser	Phe	Gln	Val	Ser	Glu	Tyr	Val	Leu	Pro	Lys	Phe	
		195					200					205				
Glu	Val	Thr	Leu	Gln	Thr	Pro	Leu	Tyr	Cys	Ser	Met	Asn	Ser	Lys	His	
	210					215					220					
Leu	Asn	Gly	Thr	Ile	Thr	Ala	Lys	Tyr	Thr	Tyr	Gly	Lys	Pro	Val	Lys	
225					230					235					240	
Gly	Asp	Val	Thr	Leu	Thr	Phe	Leu	Pro	Leu	Ser	Phe	Trp	Gly	Lys	Lys	
				245					250					255		
Lys	Asn	Ile	Thr	Lys	Thr	Phe	Lys	Ile	Asn	Gly	Ser	Ala	Asn	Phe	Ser	
		260						265					270			
Phe	Asn	Asp	Glu	Glu	Met	Lys	Asn	Val	Met	Asp	Ser	Ser	Asn	Gly	Leu	
		275					280					285				
Ser	Glu	Tyr	Leu	Asp	Leu	Ser	Ser	Pro	Gly	Pro	Val	Glu	Ile	Leu	Thr	
	290					295					300					
Thr	Val	Thr	Glu	Ser	Val	Thr	Gly	Ile	Ser	Arg	Asn	Val	Ser	Thr	Asn	
305					310					315					320	
Val	Phe	Phe	Lys	Gln	His	Asp	Tyr	Ile	Ile	Glu	Phe	Phe	Asp	Tyr	Thr	
				325					330					335		

- 32 -

Thr Val Leu Lys Pro Ser Leu Asn Phe Thr Ala Thr Val Lys Val Thr  
 340 345 350

Arg Ala Asp Gly Asn Gln Leu Thr Leu Glu Glu Arg Arg Asn Asn Val  
 355 360 365

Val Ile Thr Val Thr Gln Arg Asn Tyr Thr Glu Tyr Trp Ser Gly Ser  
 370 375 380

Asn Ser Gly Asn Gln Lys Met Glu Ala Val Gln Lys Ile Asn Tyr Thr  
 385 390 395 400

Val Pro Gln Ser Gly Thr Phe Lys Ile Glu Phe Pro Ile Leu Glu Asp  
 405 410 415

Ser Ser Glu Leu Gln Leu Lys Ala Tyr Phe Leu Gly Ser Lys Ser Ser  
 420 425 430

Met Ala Val His Ser Leu Phe Lys Ser Pro Ser Lys Thr Tyr Ile Gln  
 435 440 445

Leu Lys Thr Arg Asp Glu Asn Ile Lys Val Gly Ser Pro Phe Glu Leu  
 450 455 460

Val Val Ser Gly Asn Lys Arg Leu Lys Glu Leu Ser Tyr Met Val Val  
 465 470 475 480

Ser Arg Gly Gln Leu Val Ala Val Gly Lys Gln Asn Ser Thr Met Phe  
 485 490 495

Ser Leu Thr Pro Glu Asn Ser Trp Thr Pro Lys Ala Cys Val Ile Val  
 500 505 510

Tyr Tyr Ile Glu Asp Asp Gly Glu Ile Ile Ser Asp Val Leu Lys Ile  
 515 520 525

Pro Val Gln Leu Val Phe Lys Asn Lys Ile Lys Leu Tyr Trp Ser Lys  
 530 535 540

Val Lys Ala Glu Pro Ser Glu Lys Val Ser Leu Arg Ile Ser Val Thr  
 545 550 555 560

Gln Pro Asp Ser Ile Val Gly Ile Val Ala Val Asp Lys Ser Val Asn  
 565 570 575

Leu Met Asn Ala Ser Asn Asp Ile Thr Met Glu Asn Val Val His Glu

580	585	590
Leu Glu Leu Tyr Asn Thr Gly Tyr Tyr Leu Gly Met Phe Met Asn Ser		
595	600	605
Phe Ala Val Phe Gln Glu Cys Gly Leu Trp Val Leu Thr Asp Ala Asn		
610	615	620
Leu Thr Lys Asp Tyr Ile Asp Gly Val Tyr Asp Asn Ala Glu Tyr Ala		
625	630	635
Glu Arg Phe Met Glu Glu Asn Glu Gly His Ile Val Asp Ile His Asp		
645	650	655
Phe Ser Leu Gly Ser Ser Pro His Val Arg Lys His Phe Pro Glu Thr		
660	665	670
Trp Ile Trp Leu Asp Thr Asn Met Gly Xaa Arg Ile Tyr Gln Glu Phe		
675	680	685
Glu Val Thr Val Pro Asp Ser Ile Thr Ser Trp Val Ala Thr Gly Phe		
690	695	700
Val Ile Ser Glu Asp Leu Gly Leu Gly Leu Thr Thr Thr Pro Val Glu		
705	710	715
Leu Gln Ala Phe Gln Pro Phe Phe Ile Phe Leu Asn Leu Pro Tyr Ser		
725	730	735
Val Ile Arg Gly Glu Glu Phe Ala Leu Glu Ile Thr Ile Phe Asn Tyr		
740	745	750
Leu Lys Asp Ala Thr Glu Val Lys Val Ile Ile Glu Lys Ser Asp Lys		
755	760	765
Phe Asp Ile Leu Met Thr Ser Asn Glu Ile Asn Ala Thr Gly His Gln		
770	775	780
Gln Thr Leu Leu Val Pro Ser Glu Asp Gly Ala Thr Val Leu Phe Pro		
785	790	795
Ile Arg Pro Thr His Leu Gly Glu Ile Pro Ile Thr Val Thr Ala Leu		
805	810	815
Ser Pro Thr Ala Ser Asp Ala Val Thr Gln Met Ile Leu Val Lys Ala		
820	825	830

Glu Gly Ile Glu Lys Ser Tyr Ser Gln Ser Ile Leu Leu Asp Leu Thr  
835 840 845

Asp Asn Arg Leu Gln Ser Thr Leu Lys Thr Leu Ser Phe Ser Phe Pro  
850 855 860

Pro Asn Thr Val Thr Gly Ser Glu Arg Val Gln Ile Thr Ala Ile Gly  
865 870 875 880

Asp Val Leu Gly Pro Ser Ile Asn Gly Leu Ala Ser Leu Ile Arg Met  
885 890 895

Pro Tyr Gly Cys Gly Glu Gln Asn Met Ile Asn Phe Ala Pro Asn Ile  
900 905 910

Tyr Ile Leu Asp Tyr Leu Thr Lys Lys Lys Gln Leu Thr Asp Asn Leu  
915 920 925

Lys Glu Lys Ala Leu Ser Phe Met Arg Gln Gly Tyr Gln Arg Glu Leu  
930 935 940

Leu Tyr Gln Arg Glu Asp Gly Ser Phe Ser Ala Phe Gly Asn Tyr Asp  
945 950 955 960

Pro Ser Gly Ser Thr Trp Leu Ser Ala Phe Val Leu Arg Cys Phe Leu  
965 970 975

Glu Ala Asp Pro Tyr Ile Asp Ile Asp Gln Asn Val Leu His Arg Thr  
980 985 990

Tyr Thr Trp Leu Lys Gly His Gln Lys Ser Asn Gly Glu Phe Trp Asp  
995 1000 1005

Pro Gly Arg Val Ile His Ser Glu Leu Gln Gly Gly Asn Lys Ser  
1010 1015 1020

Pro Val Thr Leu Thr Ala Tyr Ile Val Thr Ser Leu Leu Gly Tyr  
1025 1030 1035

Arg Lys Tyr Gln Pro Asn Ile Asp Val Gln Glu Ser Ile His Phe  
1040 1045 1050

Leu Glu Ser Glu Phe Ser Arg Gly Ile Ser Asp Asn Tyr Thr Leu  
1055 1060 1065

Ala Leu Ile Thr Tyr Ala Leu Ser Ser Val Gly Ser Pro Lys Ala

1070	1075	1080
Lys Glu Ala Leu Asn Met 1085	Leu Thr Trp Arg Ala 1090	Glu Gln Glu Gly 1095
Gly Met Gln Phe Trp Val 1100	Ser Ser Glu Ser Lys 1105	Leu Ser Asp Ser 1110
Trp Gln Pro Arg Ser Leu 1115	Asp Ile Glu Val Ala 1120	Ala Tyr Ala Leu 1125
Leu Ser His Phe Leu Gln 1130	Phe Gln Thr Ser Glu 1135	Gly Ile Pro Ile 1140
Met Arg Trp Leu Ser Arg 1145	Gln Arg Asn Ser Leu 1150	Gly Gly Phe Ala 1155
Ser Thr Gln Asp Thr Thr 1160	Val Ala Leu Lys Ala 1165	Leu Ser Glu Phe 1170
Ala Ala Leu Met Asn Thr 1175	Glu Arg Thr Asn Ile 1180	Gln Val Thr Val 1185
Thr Gly Pro Ser Ser Pro 1190	Ser Pro Val Lys Phe 1195	Leu Ile Asp Thr 1200
His Asn Arg Leu Leu Leu 1205	Gln Thr Ala Glu Leu 1210	Ala Val Val Gln 1215
Pro Met Ala Val Asn Ile 1220	Ser Ala Asn Gly Phe 1225	Gly Phe Ala Ile 1230
Cys Gln Leu Asn Val Val 1235	Tyr Asn Val Lys Ala 1240	Ser Gly Ser Ser 1245
Arg Arg Arg Arg Ser Ile 1250	Gln Asn Gln Glu Ala 1255	Phe Asp Leu Asp 1260
Val Ala Val Lys Glu Asn 1265	Lys Asp Asp Leu Asn 1270	His Val Asp Leu 1275
Asn Val Cys Thr Ser Phe 1280	Ser Gly Pro Gly Arg 1285	Ser Gly Met Ala 1290
Leu Met Glu Val Asn Leu 1295	Leu Ser Gly Phe Met 1300	Val Pro Ser Glu 1305

- 36 -

Ala Ile Ser Leu Ser Glu Thr Val Lys Lys Val Glu Tyr Asp His  
1310 1315 1320

Gly Lys Leu Asn Leu Tyr Leu Asp Ser Val Asn Glu Thr Gln Phe  
1325 1330 1335

Cys Val Asn Ile Pro Ala Val Arg Asn Phe Lys Val Ser Asn Thr  
1340 1345 1350

Gln Asp Ala Ser Val Ser Ile Val Asp Tyr Tyr Glu Pro Arg Arg  
1355 1360 1365

Gln Ala Val Arg Ser Tyr Asn Ser Glu Val Lys Leu Ser Ser Cys  
1370 1375 1380

Asp Leu Cys Ser Asp Val Gln Gly Cys Arg Pro Cys Glu Asp Gly  
1385 1390 1395

Ala

<210> 9  
<211> 96  
<212> DNA  
<213> Homo sapiens

<400> 9  
attgatggtg tttatgacaa tgcagaatat gctgagaggt ttatggagga aaatgaagga 60  
catattgtag atattcatga cttttctttg ggtage 96

<210> 10  
<211> 33  
<212> PRT  
<213> homo sapiens

<400> 10

Ile Asp Gly Val Tyr Asp Asn Ala Glu Tyr Ala Glu Arg Phe Met Glu  
1 5 10 15

Glu Asn Glu Gly His Ile Val Asp Ile His Asp Phe Ser Leu Gly Ser  
20 25 30

Ser

<210> 11  
<211> 57  
<212> DNA

- 37 -

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (29)..(29)

&lt;223&gt; N = A or C

&lt;400&gt; 11

tggatttggc tagacaccaa catgggttnc aggatttacc aagaatttga agtaact

57

&lt;210&gt; 12

&lt;211&gt; 19

&lt;212&gt; PRT

&lt;213&gt; homo sapiens

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (10)..(10)

&lt;223&gt; Xaa = Ser or Tyr

&lt;400&gt; 12

Trp	Ile	Trp	Leu	Asp	Thr	Asn	Met	Gly	Xaa	Arg	Ile	Tyr	Gln	Glu	Phe
1				5					10					15	

Glu Val Thr